

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
21 October 2004 (21.10.2004)

PCT

(10) International Publication Number
WO 2004/090140 A2

(51) International Patent Classification⁷: C12N 15/82

(21) International Application Number:
PCT/EP2004/003995

(22) International Filing Date: 9 April 2004 (09.04.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03076044.1 9 April 2003 (09.04.2003) EP
60/496,688 21 August 2003 (21.08.2003) US

(71) Applicant (for all designated States except US): BAYER
BIOSCIENCE N.V. [BE/BE]; Technologiepark 38,
B-9052 Gent (BE).

(72) Inventor; and

(75) Inventor/Applicant (for US only): DE BLOCK, Marc
[BE/BE]; Abrikozenstraat 26, B-9820 Merelbeke (BE).

(74) Common Representative: BAYER BIOSCIENCE N.V.;
Technologiepark 38, B-9052 Gent (BE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.


(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW),
Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Euro-
pean (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR,
GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to the identity of the inventor (Rule 4.17(i)) for the fol-
lowing designations AE, AG, AL, AM, AT, AU, AZ, BA, BB,
BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE,
DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD,
SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ,

[Continued on next page]

(54) Title: METHODS AND MEANS FOR INCREASING THE TOLERANCE OF PLANTS TO STRESS CONDITIONS



(57) Abstract: Methods and means are provided to increase the tolerance of plants to abiotic stress or adverse growing conditions, including drought, high light intensities, high temperatures, nutrient limitations and the like by reducing the activity of endogenous PARG proteins in plants.



VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, SD,

SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations
- of inventorship (Rule 4.17(iv)) for US only

Published:

- without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

10/552552

Methods and means for increasing the tolerance of plants to stress conditions.

Field of the invention.

[1] The present invention relates to the use of poly (ADP-ribose) glycohydrolases in plants to increase the tolerance of plants to adverse growing conditions, including drought, high light intensities, high temperatures, nutrient limitations and the like. Methods and means are provided to produce plants that are tolerant to abiotic stress conditions.

Background to the invention

[2] . Frequently, abiotic stress will lead either directly or indirectly to damage of the DNA of the cells of the plants exposed to the adverse conditions. Genomic damage, if left unrepaired, can lead to cell death. Tolerance to stress conditions exhibited by plants is the result of the ability of the plant cells exposed to the adverse conditions to reduce and/or repair the damage, and to survive.

[3] Plant cells, like other eukaryotic cells, have evolved an elaborate DNA repair system. The activation of poly(ADP-ribose) polymerase (PARP) by DNA strand breaks is often one of the first cellular responses to DNA damage. PARP catalyzes the post-translational modification of proteins by adding successively molecules of ADP-ribose, obtained from the conversion of nicotineamide dinucleotide (NAD), to form multibranched polymers containing up to 200 ADP-ribose residues (about 40 residues in plants). The dependence of poly(ADP-ribose) synthesis on DNA strand breaks, and the presence of PARP in multiprotein complexes further containing key effectors of DNA repair, replication and transcription reactions, strongly suggests that this posttranslational modification is involved in metabolism of nucleic acids, and DNA repair. There are also indications that poly (ADP -ribose) synthesis is involved in regulation of cell cycle and cell death.

[4] Poly (ADP-ribosylation) of proteins is transient in living cells. The poly (ADP-ribose) polymers are rapidly turned over, being converted to free ADP-ribose by the exoglycosidase and endoglycosidase activity of poly (ADP-ribose) glycohydrolase (PARG; E.C.3.2.1.143). The most proximal unit of ADP-ribose on the protein acceptor is hydrolyzed by the action of another enzyme (ADP-ribosyl protein lyase).

[5] In addition to this positive (DNA-repair associated) effect of PARP on cell survival, there is also a negative effect of PARP. The process of activating PARP upon DNA damage is associated with a rapid lowering of NAD⁺ levels, since each ADP-ribose unit transferred by PARP consumes one molecule of NAD⁺. NAD⁺ depletion in turn results in ATP depletion, because NAD⁺ resynthesis requires at least (depending on the biosynthesis pathway) three molecules of ATP per molecule of NAD⁺. Furthermore, NAD⁺ depletion blocks glyceraldehyde -3-phosphate dehydrogenase activity, which is required to resynthesize ATP during glycolysis. Finally, NAD⁺ is a key carrier of electrons needed to generate ATP via electron transport and oxidative phosphorylation.

[6] The physiological consequence of NAD⁺ and ATP depletion has been established in the context of DNA-damage induced cell death. It has been shown that the completion of apoptosis is absolutely dependent on the presence of ATP and that, in the absence of this nucleotide, the type of cellular demise switches from apoptosis to necrosis. Since the cellular lysis associated with necrosis generates further damage to neighboring cells it is preferable for multicellular organisms to favor apoptotic cell death rather than necrosis.

[7] It is thus very important to consider the delicate balance of positive and negative effects of the poly (ADP-ribosylation) on the potential of a cell to survive DNA damage.

[8] WO 00/04173 describes methods to modulate programmed cell death (PCD) in eukaryotic cells and organisms, particularly plant cells and plants, by

introducing of "PCD modulating chimeric genes" influencing the expression and/or apparent activity of endogenous poly-(ADP-ribose) polymerase (PARP) genes. Programmed cell death may be inhibited or provoked. The invention particularly relates to the use of nucleotide sequences encoding proteins with PARP activity for modulating PCD, for enhancing growth rate or for producing stress tolerant cells and organisms.

[9] PARP encoding genes have been identified in a number of animals such as *Rattus norvegicus* (Accession numbers: NM_031339, NW_043030, AB019366,), *Mus musculus* (Accession numbers: NT_039598, NM_003631, AF079557), *Homo sapiens* (Accession numbers: NT_017696; NM_003631, AF005043), *Bos taurus* (Accession numbers: NM_174138, U78975) *Drosophila melanogaster* (Accession number: AF079556)

[10] In plants, a poly(ADP-ribose) glycohydrolase has been identified by map-based cloning of the wild-type gene inactivated in a mutant affected in clock-controlled transcription of genes in *Arabidopsis* and in photoperiod dependent transition from vegetative growth to flowering (*tef*). The nucleotide sequence of the gene can be obtained from nucleotide databases under the accession number AF394690 (Panda et al., 2002 Dev. Cell. 3, 51-61).

Summary of the invention

[11] The invention provides a method to produce a plant tolerant to stress conditions comprising the steps of providing plant cells with a chimeric gene to create transgenic plant cells, wherein the chimeric gene comprises the following operably linked DNA fragments: a plant-expressible promoter; a DNA region, which when transcribed yields an ParG inhibitory RNA molecule; and a 3' end region involved in transcription termination and polyadenylation. A population of transgenic plant lines is regenerated from the transgenic plant cell; and a stress tolerant plant line is identified within the population of transgenic plant lines. The ParG inhibitory RNA molecule may comprise a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG

gene). The ParG inhibitory RNA molecule may also comprise a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG gene). In yet another embodiment, the parG inhibitory RNA may comprise a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell, wherein the sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides. The chimeric gene may further comprise a DNA region encoding a self-splicing ribozyme between said DNA region coding for parG inhibitory RNA molecule and the 3' end region. Stress conditions may be selected from heat, drought, nutrient depletion, oxidative stress or high light conditions.

[12] In another embodiment of the invention, a method is provided to produce a plant tolerant to stress conditions comprising the steps of: isolating a DNA fragment of at least 100 bp comprising a part of the parG encoding gene of the plant of interest; producing a chimeric gene by operably linking a plant expressible promoter region to the isolated DNA fragment comprising part of the parG encoding gene of the plant in direct orientation compared to the promoter region; and to the isolated DNA fragment comprising part of the parG encoding gene of said plant in inverted orientation compared to the promoter region, and a 3' end region involved in transcription termination and polyadenylation. These chimeric genes are then provided to plant cells to create transgenic plant cells. A population of transgenic plant lines is regenerated from the transgenic plant cells; and a stress tolerant plant line is identified within the population of transgenic plant lines. The invention also relates to stress tolerant plant cells and plants obtained by this process.

[13] In yet another embodiment of the invention, a method is provided to produce a plant tolerant to stress conditions comprising the steps of providing plant cells with a chimeric gene to create transgenic plant cells, comprising a

DNA region, which when transcribed yields an ParG inhibitory RNA molecule, whereby the DNA region comprises a nucleotide sequence of at least 21 to 100 nucleotides of a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID No 1, 2 or 16 or at least 21 to 100 nucleotides of a nucleotide sequence of SEQ ID 3, 4, 15 or 23 operably linked to a plant-expressible promoter and a 3' end region involved in transcription termination and polyadenylation; regenerating a population of transgenic plant lines from said transgenic plant cell; and identifying a stress tolerant plant line within the population of transgenic plant lines.

[14] The invention also provides DNA molecules comprising a plant-expressible promoter, operably linked to a DNA region, which when transcribed yields an ParG inhibitory RNA molecule, and to a 3' end region involved in transcription termination and polyadenylation. The ParG inhibitory RNA molecule may comprise a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG gene). The ParG inhibitory RNA molecule may also comprise a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell (the endogenous ParG gene). In yet another embodiment, the parG inhibitory RNA may comprise a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in the plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in the plant cell, wherein the sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides. The chimeric gene may further comprise a DNA region encoding a self-splicing ribozyme between said DNA region coding for parG inhibitory RNA molecule and the 3' end region. The chimeric gene may also comprise a nucleotide sequence of at least 21 to 100 nucleotides of a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID No 1, 2 or 16 or at least 21 to 100 nucleotides of a nucleotide sequence of SEQ ID 3, 4, 15 or 23.

[15] In yet another embodiment, the invention relates to plant cell comprising the DNA molecule of the invention and plants consisting essentially of such plant cells, as well as to processes for producing stress tolerant plants, comprising the step of further crossing such plants with another plant. Seeds and propagating material of such plants comprising the chimeric genes of the invention are also provided.

[16] The invention also relates to a method for obtaining stress tolerant plants comprising the steps of subjecting a plant cell line or a plant or plant line, to mutagenesis; identifying those plant cells or plants that have a mutation in an endogenous ParG gene; subjecting the identified plant cells or plants to stress conditions and identifying plant cells or plants that tolerate said stress conditions better than control plants. Alternatively, plant cells or plants may be selected for resistance to ParG inhibitors and further treated as described in this paragraph.

[17] The invention further relates to a stress tolerant plant cell or plant having a mutation in the endogenous ParG gene.

Brief description of the figures.

[18] Figure 1. Schematic representation of the poly-ADP ribose polymerization / depolymerization cycle by the action of PARP/PARG in a eukaryotic cell.

[19] Figure 2. Diagram of the NAD⁺ and ATP content of *Arabidopsis* lines under high light stress. Dark boxes represent NAD content under high light conditions expressed as percentage of the value for NAD content determined under low light conditions. Light boxes represent ATP content under high light conditions expressed as percentage of the value for ATP content determined under low light conditions.

[20] Figure 3. Diagram of the NAD⁺ and ATP content of corn lines under nitrogen depletion stress. Dark boxes represent NAD content while light boxes represent ATP content.

Detailed description of preferred embodiments

[21] The invention is based, on the one hand, on the demonstration that cells from stress resistant plant lines comprising a chimeric gene reducing the PARP gene expression, exhibited a higher NAD/ATP content under adverse conditions than cells from untransformed plant lines. On the other hand, it has been observed that silencing of the expression of PARG encoding gene in tobacco using a transient silencing RNA vector based on satellite viruses resulted in a similar phenotype as that observed for silencing of PARP encoding gene using the same silencing system. Furthermore, silencing the expression of PARG encoding gene in plants, such as Arabidopsis and tobacco, resulted in plants that were more resistant to stress conditions, such as e.g. those imposed by high light conditions.

[22] Although not intending to limit the invention to a specific mode of action, it is expected that silencing of PARG gene expression results in a similar phenotype as silencing of PARP gene expression for the following reasons. As can be seen from Figure 1, polymerization of ADP ribose catalyzed by PARP, consuming NAD, is followed by depolymerization of poly ADP ribose, catalyzed by PARG. Poly ADP ribosylation of the PARP protein itself results in inactivation of the PARP protein. The speed at which the ADP ribose polymerization / depolymerization cycle occurs in plant cells, leading to NAD depletion and consequently ATP depletion, can be slowed down or stopped by reduction of the PARP gene expression or of the enzymatic activity of PARP. As a result, plant cells, and plants comprising such cells are more resistant to adverse conditions. The data provided here indicate that a similar effect can be obtained through slowing down or stopping the cycle by reduction of the PARG gene expression or PARG activity.

[23] The invention relates to reduction of plant cell death in response to adverse environmental conditions, and consequently to enhanced stress resistance, by altering the level of expression of ParG genes, or by altering the activity or the apparent activity of PARG proteins in that plant cell. Conveniently, the level of expression of ParG genes may be controlled genetically by introduction of chimeric genes altering the expression of ParG genes, or by altering the endogenous PARG encoding genes, including the expression signals.

[24] In one embodiment of the invention, a method for producing plants tolerant to stress conditions or adverse growing conditions is provided comprising the steps of:

- providing plant cells with a chimeric gene to create transgenic plant cells, wherein the chimeric gene comprises the following operably linked DNA fragments:
 - a plant-expressible promoter;
 - a DNA region, which when transcribed yields a ParG inhibitory RNA molecule;
 - a 3' end region involved in transcription termination and polyadenylation;
- regenerating a population of transgenic plant lines from said transgenic plant cell; and
- identifying a stress tolerant plant line within said population of transgenic plant lines.

[25] As used herein "a stress tolerant plant" or "a plant tolerant to stress conditions or adverse growing conditions" is a plant (particularly a plant obtained according to the methods of the invention), which, when subjected to adverse growing conditions for a period of time, such as but not limited to drought, high temperatures, limited supply of nutrients (particularly nitrogen), high light intensities, grows better than a control plant not treated according to the methods of the invention. This will usually be apparent from the general appearance of the plants and may be measured e.g., by increased biomass production, continued vegetative growth under adverse conditions or higher

seed yield. Stress tolerant plant have a broader growth spectrum, i.e. they are able to withstand a broader range of climatological and other abiotic changes, without yield penalty. Biochemically, stress tolerance may be apparent as the higher NAD^+ -NADH /ATP content and lower production of reactive oxygen species of stress tolerant plants compared to control plants under stress condition. Stress tolerance may also be apparent as the higher chlorophyll content, higher photosynthesis and lower chlorophyll fluorescence under stress conditions in stress tolerant plants compared to control plants under the same conditions.

[26] It will be clear that it is also not required that the plant be grown continuously under the adverse conditions for the stress tolerance to become apparent. Usually, the difference in stress tolerance between a plant or plant cell according to the invention and a control plant or plant cell will become apparent even when only a relatively short period of adverse conditions is encountered during growth.

[27] As used herein, a "ParG inhibitory RNA molecule" is an RNA molecule that is capable of decreasing the expression of the endogenous PARG encoding genes of a plant cell, preferably through post-transcriptional silencing. It will be clear that even when a ParG inhibitory RNA molecule decreases the expression of a PARG encoding gene through post-transcriptional silencing, such an RNA molecule may also exert other functions within a cell, such as e.g. guiding DNA methylation of the endogenous ParG gene, again ultimately leading to decreased expression of the PARG encoding gene. Also, expression of the endogenous PARG encoding genes of a plant cell may be reduced by transcriptional silencing, e.g., by using RNAi or dsRNA targeted against the promoter region of the endogenous ParG gene.

[28] As used herein, a "PARG encoding gene" or a "ParG gene" is a gene capable of encoding a PARG (poly ADP ribose glycohydrolase) protein, wherein the PARG protein catalyzes the depolymerization of poly ADP-ribose,

by releasing free ADP ribose units either by endoglycolytic or exoglycolytic action.

[29] PARG encoding genes may comprise a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID No 1 (*Arabidopsis thaliana*) or of SEQ ID No 2 (*Solanum tuberosum*) or of SEQ ID No 16 (*Oryza sativa*) or parts thereof, such as a DNA fragment comprising the nucleotide sequence of SEQ ID No. 3 or SEQ ID 4 or SEQ ID No 15. or SEQ ID 23 (*Zea mays*).

[30] However, it will be clear that the skilled person can isolate variant DNA sequences from other plant species, by hybridization with a probe derived from the above mentioned PARG encoding genes from plant species, or even with a probe derived from the above mentioned PARG encoding genes from animal species. To this end, the probes should preferably have a nucleotide sequence comprising at least 40 consecutive nucleotides from the coding region of those mentioned PARG encoding genes sequences, preferably from the coding region of SEQ ID No 3 or SEQ ID No 4. The probes may however comprise longer regions of nucleotide sequences derived from the ParG genes, such as about 50, 60, 75, 100, 200 or 500 consecutive nucleotides from any of the mentioned ParG genes. Preferably, the probe should comprise a nucleotide sequence coding for one of the highly conserved regions of the catalytic domain, which have been identified by aligning the different PARG proteins from animals. These regions are also present in the identified PARG protein from *Arabidopsis thaliana* and comprise the amino acid sequence LXVDFANXXXGGG (corresponding to SEQ ID No 1 from the amino acid at position 252 to the amino acid at position 264; X may be any amino acid) LXVDFANXXXGGGXXXGXVQEEIRF (corresponding to SEQ ID No 1 from the amino acid at position 252 to the amino acid at position 277) or LXVDFANXXXGGGXXXGXVQEEIRFXXXPE (corresponding to SEQ ID No 1 from the amino acid at position 252 to the amino acid at position 282), TGXWGCXFXGD (corresponding to SEQ ID No 1 from the amino acid at position 449 to the amino acid at position 460) or TGXWGCXGAFXGDXXLKXXXQ (corresponding to SEQ ID No 1 from the

amino acid at position 449 to the amino acid at position 468). Other conserved regions have the amino acid sequence DXXXRXXXXAIDA (corresponding to SEQ ID No 1 from the amino acid at position 335 to the amino acid at position 344) or REXXKAXXGF (corresponding to SEQ ID No 1 from the amino acid at position 360 to the amino acid at position 369) or GXXXXSXYTGY (corresponding to SEQ ID No 1 from the amino acid at position 303 to the amino acid at position 313). Hybridization should preferably be under stringent conditions.

[31] "Stringent hybridization conditions" as used herein mean that hybridization will generally occur if there is at least 95% and preferably at least 97% sequence identity between the probe and the target sequence. Examples of stringent hybridization conditions are overnight incubation in a solution comprising 50% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5x Denhardt's solution, 10% dextran sulfate, and 20 µg/ml denatured, sheared carrier DNA such as salmon sperm DNA, followed by washing the hybridization support in 0.1 x SSC at approximately 65 °C, e.g. for about 10 min (twice). Other hybridization and wash conditions are well known and are exemplified in Sambrook et al, Molecular Cloning: A Laboratory Manual, Second Edition, Cold Spring Harbor, NY (1989), particularly chapter 11.

[32] Alternatively, ParG encoding genes or parts thereof may also be isolated by PCR based techniques, using as primers oligonucleotides comprising at least 20 consecutive nucleotides from a nucleotide sequence of the mentioned PARG encoding genes or the complement thereof. Such primers may comprise a nucleotide sequence encoding a conserved region, as mentioned above, or be complementary to such a nucleotide sequence. Oligonucleotides which may be used for that purpose may comprise the nucleotide sequence of either or SEQ ID No.5, SEQ ID No 6., SEQ ID No. 7 or SEQ ID No. 8. Oligonucleotides which may be used may also be degenerate, such as the oligonucleotide primers of SEQ ID No 17, SEQ ID No 18, SEQ ID No 19; SEQ ID No 20, SEQ ID No 21 or SEQ ID No 22.

[33] Specific PCR fragments from ParG genes may e.g., be obtained by using combinations of the oligonucleotides having the nucleotide sequence of SEQ ID No. 5 and SEQ ID No 6 using e.g., *Arabidopsis* genomic DNA or cDNA as a template DNA, or by using combinations of the oligonucleotides having the nucleotide sequence of SEQ ID No. 7 and SEQ ID No 8 using e.g., potato genomic DNA or cDNA as a template DNA, under stringent annealing conditions.

[34] The isolated sequences may encode a functional PARG protein or a part thereof. Preferably the isolated sequences should comprise a nucleotide sequence coding for one or more of the highly conserved regions from the catalytic domain of PARG proteins as mentioned elsewhere.

[35] However, for the purpose of the invention is not required that the isolated sequences encode a functional ParG protein nor that a complete coding region is isolated. Indeed, all that is required for the invention is that a chimeric gene can be designed or produced, based on the identified or isolated sequence of the endogenous ParG gene from a plant, which is capable of producing a ParG inhibitory RNA. Several alternative methods are available to produce such a ParG inhibitory RNA molecule.

[36] In one embodiment, the ParG inhibitory RNA molecule encoding chimeric gene is based on the so-called antisense technology. In other words, the coding region of the chimeric gene comprises a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the endogenous ParG gene of the plant cell or plant, the expression of which is targeted to be reduced. Such a chimeric gene may be conveniently constructed by operably linking a DNA fragment comprising at least 20 nucleotides from the isolated or identified ParG gene, or part of such a gene, in inverse orientation, to a plant expressible promoter and 3'end formation region involved in transcription termination and polyadenylation. It will be immediately clear that there is no need to know the exact nucleotide sequence or complete nucleotide sequence of such a DNA fragment from an isolated ParG gene.

[37] In another embodiment the ParG inhibitory RNA molecule encoding chimeric gene is based on the so-called co-suppression technology. In other words, the coding region of the chimeric gene comprises a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the endogenous ParG gene of the plant cell or plant, the expression of which is targeted to be reduced. Such a chimeric gene may be conveniently constructed by operably linking a DNA fragment comprising at least 20 nucleotides from the isolated or identified ParG gene, or part of such a gene, in direct orientation, to a plant expressible promoter and 3'end formation region involved in transcription termination and polyadenylation. Again it is not required to know the exact nucleotide sequence of the used DNA fragment from the isolated ParG gene.

[38] The efficiency of the above mentioned chimeric genes in reducing the expression of the endogenous ParG gene may be further enhanced by inclusion of DNA elements which result in the expression of aberrant, unpolyadenylated ParG inhibitory RNA molecules. One such DNA element suitable for that purpose is a DNA region encoding a self-splicing ribozyme, as described in WO 00/01133.

[39] The efficiency of the above mentioned chimeric genes in reducing the expression of the endogenous ParG gene of a plant cell may also be further enhanced by including into one plant cell simultaneously a chimeric gene as herein described encoding an antisense ParG inhibitory RNA molecule and a chimeric gene as herein described encoding a sense ParG inhibitory RNA molecule, wherein said antisense and sense ParG inhibitory RNA molecules are capable of forming a double stranded RNA region by base pairing between the mentioned at least 20 consecutive nucleotides, as described in WO 99/53050.

[40] As further described in WO 99/53050, the sense and antisense ParG inhibitory RNA regions, capable of forming a double stranded RNA region may be present in one RNA molecule, preferably separated by a spacer region.

The spacer region may comprise an intron sequence. Such a chimeric gene may be conveniently constructed by operably linking a DNA fragment comprising at least 20 nucleotides from the isolated or identified endogenous ParG gene, the expression of which is targeted to be reduced, in an inverted repeat, to a plant expressible promoter and 3' end formation region involved in transcription termination and polyadenylation. To achieve the construction of such a chimeric gene, use can be made of the vectors described in WO 02/059294

[41] An embodiment of the invention thus concerns a method for obtaining a stress tolerant plant line comprising the steps of

- providing plant cells with a chimeric gene to create transgenic plant cells, wherein the chimeric gene comprises the following operably linked DNA fragments:

- a plant-expressible promoter;
- a DNA region, which when transcribed yields a ParG inhibitory RNA molecule comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell; or
- a DNA region, which when transcribed yields a ParG inhibitory RNA molecule comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell; or
- a DNA region, which when transcribed yields a ParG inhibitory RNA molecule comprising a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell, wherein said sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides.

- a 3' end region involved in transcription termination and polyadenylation;
- regenerating a population of transgenic plant lines from said transgenic plant cell; and
- identifying a stress tolerant plant line within said population of transgenic plant lines.

[42] As used herein "comprising" is to be interpreted as specifying the presence of the stated features, integers, steps or components as referred to, but does not preclude the presence or addition of one or more features, integers, steps or components, or groups thereof. Thus, e.g., a nucleic acid or protein comprising a sequence of nucleotides or amino acids, may comprise more nucleotides or amino acids than the actually cited ones, i.e., be embedded in a larger nucleic acid or protein. A chimeric gene comprising a DNA region which is functionally or structurally defined, may comprise additional DNA regions etc.

[43] It will thus be clear that the minimum nucleotide sequence of the antisense or sense RNA region of about 20 nt of the ParG coding region may be comprised within a larger RNA molecule, varying in size from 20 nt to a length equal to the size of the target gene.

[44] The mentioned antisense or sense nucleotide regions may thus be about from about 21 nt to about 5000 nt long, such as 21nt, 40 nt, 50 nt, 100nt, 200 nt, 300nt, 500nt, 1000 nt, 2000 nt or even about 5000 nt or larger in length.

[45] Moreover, it is not required for the purpose of the invention that the nucleotide sequence of the used inhibitory ParG RNA molecule or the encoding region of the chimeric gene, is completely identical or complementary to the endogenous ParG gene the expression of which is targeted to be reduced in the plant cell. The longer the sequence, the less stringent the requirement for the overall sequence identity is. Thus, the sense or antisense regions may have an overall sequence identity of about 40 % or

50% or 60 % or 70% or 80% or 90 % or 100% to the nucleotide sequence of the endogenous ParG gene or the complement thereof. However, as mentioned antisense or sense regions should comprise a nucleotide sequence of 20 consecutive nucleotides having about 100% sequence identity to the nucleotide sequence of the endogenous ParG gene. Preferably the stretch of about 100 % sequence identity should be about 50, 75 or 100 nt.

[46] For the purpose of this invention, the "sequence identity" of two related nucleotide sequences, expressed as a percentage, refers to the number of positions in the two optimally aligned sequences which have identical residues (x100) divided by the number of positions compared. A gap, i.e. a position in an alignment where a residue is present in one sequence but not in the other is regarded as a position with non-identical residues. The alignment of the two sequences is performed by the Needleman and Wunsch algorithm (Needleman and Wunsch 1970) Computer-assisted sequence alignment, can be conveniently performed using standard software program such as GAP which is part of the Wisconsin Package Version 10.1 (Genetics Computer Group, Madison, Wisconsin, USA) using the default scoring matrix with a gap creation penalty of 50 and a gap extension penalty of 3.

[47] It will be clear that whenever nucleotide sequences of RNA molecules are defined by reference to nucleotide sequence of corresponding DNA molecules, the thymine (T) in the nucleotide sequence should be replaced by uracil (U). Whether reference is made to RNA or DNA molecules will be clear from the context of the application.

[48] It will also be clear that chimeric genes capable of producing inhibitory ParG genes for a particular ParG gene in a particular plant variety or plant species, may also be used to inhibit ParG gene expression in other plant varieties or plant species. Indeed, when sufficient homology exists between the ParG inhibitory RNA region and the ParG gene, or when the ParG genes share the same stretch of 19 nucleotides, expression of those other genes will also be down-regulated.

[49] In view of the potential role of ParG in nucleic acid metabolism, it may be advantageous that the expression of the endogenous ParG gene by the ParG inhibitory RNA is not completely inhibited. Downregulating the expression of a particular gene by gene silencing through the introduction of a chimeric gene encoding ParG inhibitory RNA will result in a population of different transgenic lines, exhibiting a distribution of different degrees of silencing of the ParG gene. The population will thus contain individual transgenic plant lines, wherein the endogenous ParG gene is silenced to the required degree of silencing. A person skilled in the art can easily identify such plant lines, e.g. by subjecting the plant lines to a particular adverse condition, such a high light intensity, oxidative stress, drought, heat etc. and selecting those plants which perform satisfactory and survive best the treatment.

[50] As used herein, the term "promoter" denotes any DNA which is recognized and bound (directly or indirectly) by a DNA-dependent RNA-polymerase during initiation of transcription. A promoter includes the transcription initiation site, and binding sites for transcription initiation factors and RNA polymerase, and can comprise various other sites (e.g., enhancers), at which gene expression regulatory proteins may bind.

[51] The term "regulatory region", as used herein, means any DNA, that is involved in driving transcription and controlling (i.e., regulating) the timing and level of transcription of a given DNA sequence, such as a DNA coding for a protein or polypeptide. For example, a 5' regulatory region (or "promoter region") is a DNA sequence located upstream (i.e., 5') of a coding sequence and which comprises the promoter and the 5'-untranslated leader sequence. A 3' regulatory region is a DNA sequence located downstream (i.e., 3') of the coding sequence and which comprises suitable transcription termination (and/or regulation) signals, including one or more polyadenylation signals.

[52] In one embodiment of the invention the promoter is a constitutive promoter. In another embodiment of the invention, the promoter activity is enhanced by external or internal stimuli (inducible promoter), such as but not

limited to hormones, chemical compounds, mechanical impulses, abiotic or biotic stress conditions. The activity of the promoter may also regulated in a temporal or spatial manner (tissue-specific promoters; developmentally regulated promoters).

[53] For the purpose of the invention, the promoter is a plant-expressible promoter. As used herein, the term "plant-expressible promoter" means a DNA sequence which is capable of controlling (initiating) transcription in a plant cell. This includes any promoter of plant origin, but also any promoter of non-plant origin which is capable of directing transcription in a plant cell, i.e., certain promoters of viral or bacterial origin such as the CaMV35S (Hapster et al., 1988), the subterranean clover virus promoter No 4 or No 7 (WO9606932), or T-DNA gene promoters but also tissue-specific or organ-specific promoters including but not limited to seed-specific promoters (e.g., WO89/03887), organ-primordia specific promoters (An et al., 1996), stem-specific promoters (Keller et al., 1988), leaf specific promoters (Hudspeth et al., 1989), mesophyl-specific promoters (such as the light-inducible Rubisco promoters), root-specific promoters (Keller et al., 1989), tuber-specific promoters (Keil et al., 1989), vascular tissue specific promoters (Peleman et al., 1989), stamen-selective promoters (WO 89/10396, WO 92/13956), dehiscence zone specific promoters (WO 97/13865) and the like.

[54] Methods for the introduction of chimeric genes into plants are well known in the art and include *Agrobacterium*-mediated transformation, particle gun delivery, microinjection, electroporation of intact cells, polyethyleneglycol-mediated protoplast transformation, electroporation of protoplasts, liposome-mediated transformation, silicon-whiskers mediated transformation etc. The transformed cells obtained in this way may then be regenerated into mature fertile plants.

[55] The transgenic plant cells and plant lines according to the invention may further comprise chimeric genes which will reduce the expression of PARP genes as described in WO 00/04173. These further chimeric genes may be introduced e.g. by crossing the transgenic plant lines of the current

invention with transgenic plants containing PARP gene expression reducing chimeric genes. Transgenic plant cells or plant lines may also be obtained by introducing or transforming the chimeric genes of the invention into transgenic plant cells comprising the PARP gene expression reducing chimeric genes or vice versa. Alternatively, the PARP and PARG inhibitory RNA regions may be encoded by one chimeric gene and transcribed as one RNA molecule.

[56] The chimeric genes of the invention (or the inhibitory RNA molecules corresponding thereto) may also be introduced into plant cells in a transient manner, e.g. using the viral vectors, such as viral RNA vectors as described in WO 00/63397 or WO 02/13964.

[57] Having read this specification, it will be immediately clear to the skilled artisan, that mutant plant cells and plant lines, wherein the PARG activity is reduced may be used to the same effect as the transgenic plant cells and plant lines described herein. Mutants in ParG gene of a plant cell or plant may be easily identified using screening methods known in the art, whereby chemical mutagenesis, such as e.g., EMS mutagenesis, is combined with sensitive detection methods (such as e.g., denaturing HPLC). An example of such a technique is the so-called "Targeted Induced Local Lesions in Genomes" method as described in McCallum et al, Plant Physiology 123 439-442 or WO 01/75167. However, other methods to detect mutations in particular genome regions or even alleles, are also available and include screening of libraries of existing or newly generated insertion mutant plant lines, whereby pools of genomic DNA of these mutant plant lines are subjected to PCR amplification using primers specific for the inserted DNA fragment and primers specific for the genomic region or allele, wherein the insertion is expected (see e.g. Maes et al., 1999, Trends in Plant Science, 4, pp 90-96).

[58] Plant cell lines and plant lines may also be subjected to mutagenesis by selection for resistance to ParG inhibitors, such as gallotannines . (Ying, et al. (2001). Proc. Natl. Acad. Sci. USA 98(21), 12227-12232; Ying, W., Swanson, R.A. (2000). NeuroReport 11(7), 1385-1388.

[59] Thus, methods are available in the art to identify plant cells and plant lines comprising a mutation in the ParG gene. This population of mutant cells or plant lines can then be subjected to different abiotic stresses, and their phenotype or survival can be easily determined. Additionally, the NAD and/or the ATP content of the stressed cells can be determined and compared to results of such determinations of unstressed cells. In stress tolerant cells, the reduction of NAD content under stress conditions should when compared with unstressed cells, should be lower than for corresponding control cells.

[60] It is also an object of the invention to provide plant cells and plants containing the chimeric genes or the RNA molecules according to the invention. Gametes, seeds, embryos, either zygotic or somatic, progeny or hybrids of plants comprising the chimeric genes of the present invention, which are produced by traditional breeding methods are also included within the scope of the present invention.

[61] The plants obtained by the methods described herein may be further crossed by traditional breeding techniques with other plants to obtain stress tolerant progeny plants comprising the chimeric genes of the present invention.

[62] The methods and means described herein are believed to be suitable for all plant cells and plants, both dicotyledonous and monocotyledonous plant cells and plants including but not limited to cotton, Brassica vegetables, oilseed rape, wheat, corn or maize, barley, alfalfa, peanuts, sunflowers, rice, oats, sugarcane, soybean, turf grasses, barley, rye, sorghum, sugar cane, vegetables (including chicory, lettuce, tomato, zucchini, bell pepper, eggplant, cucumber, melon, onion, leek), tobacco, potato, sugarbeet, papaya, pineapple, mango, *Arabidopsis thaliana*, but also plants used in horticulture, floriculture or forestry (poplar, fir, eucalyptus etc.).

[63] The following non-limiting Examples describe method and means for increasing stress tolerance in plants according to the invention.

[64] Unless stated otherwise in the Examples, all recombinant DNA techniques are carried out according to standard protocols as described in Sambrook et al. (1989) *Molecular Cloning: A Laboratory Manual*, Second Edition, Cold Spring Harbor Laboratory Press, NY and in Volumes 1 and 2 of Ausubel et al. (1994) *Current Protocols in Molecular Biology*, Current Protocols, USA. Standard materials and methods for plant molecular work are described in *Plant Molecular Biology Labfax* (1993) by R.D.D. Croy, jointly published by BIOS Scientific Publications Ltd (UK) and Blackwell Scientific Publications, UK. Other references for standard molecular biology techniques include Sambrook and Russell (2001) *Molecular Cloning: A Laboratory Manual*, Third Edition, Cold Spring Harbor Laboratory Press, NY, Volumes I and II of Brown (1998) *Molecular Biology LabFax*, Second Edition, Academic Press (UK). Standard materials and methods for polymerase chain reactions can be found in Dieffenbach and Dveksler (1995) *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, and in McPherson et al. (2000) *PCR - Basics: From Background to Bench*, First Edition, Springer Verlag, Germany.

[65] Throughout the description and Examples, reference is made to the following sequences:

[66] SEQ ID N°1: amino acid sequence of the ParG protein from *Arabidopsis thaliana*.

[67] SEQ ID N°2: amino acid sequence of part of the ParG protein from *Solanum tuberosum*.

[68] SEQ ID N°3: nucleotide sequence encoding the ParG protein from *Arabidopsis thaliana*.

[69] SEQ ID N°4: nucleotide sequence encoding the part of the ParG protein from *Solanum tuberosum*.

[70] SEQ ID N°5: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.

[71] SEQ ID N°6: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.

- [72] SEQ ID N°7: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [73] SEQ ID N°8: nucleotide sequence of an oligonucleotide primer suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [74] SEQ ID N°9: nucleotide sequence of the T-DNA vector containing the ParG expression reducing chimeric gene based on the *Arabidopsis* ParG gene sequence.
- [75] SEQ ID N°10: amino acid sequence of conserved sequence 1 of PARG proteins.
- [76] SEQ ID N°11: amino acid sequence of conserved sequence 2 of PARG proteins.
- [77] SEQ ID N°12: amino acid sequence of conserved sequence 3 of PARG proteins.
- [78] SEQ ID N°13: amino acid sequence of conserved sequence 4 of PARG proteins.
- [79] SEQ ID N°14: amino acid sequence of conserved sequence 5 of PARG proteins.
- [80] SEQ ID N°15: nucleotide sequence of the ParG protein from *Oryza sativa*.
- [81] SEQ ID N°16: amino acid sequence of the ParG protein from *Oryza sativa*.
- [82] SEQ ID N° 17: nucleotide sequence of an oligonucleotide primer PG1 suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [83] SEQ ID N° 18: nucleotide sequence of an oligonucleotide primer PG2 suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [84] SEQ ID N° 19: nucleotide sequence of an oligonucleotide primer PG3 suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [85] SEQ ID N° 20: nucleotide sequence of an oligonucleotide primer PG4 suitable for PCR amplification of part of a ParG protein encoding DNA fragment.

- [86] SEQ ID N° 21: nucleotide sequence of an oligonucleotide primer PG5 suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [87] SEQ ID N° 22: nucleotide sequence of an oligonucleotide primer PG6 suitable for PCR amplification of part of a ParG protein encoding DNA fragment.
- [88] SEQ ID N°23: nucleotide sequence encoding a ParG protein from *Zea mays*.
- [89] SEQ ID N°24: nucleotide sequence of a T-DNA vector comprising a chimeric gene capable of reducing PARG expression
- [90] SEQ ID N°25: nucleotide sequence of a T-DNA vector comprising a chimeric gene capable of reducing PARG expression

[91] Examples

Example 1. Analysis of the influence of stress on energy production efficiency of transgenic stress tolerant plant lines containing PARP gene expression reducing chimeric genes.

[92] Hypocotyls of transgenic *Brassica napus* plants comprising PARP gene expression reducing chimeric genes as described in WO 00/04173 were cultivated for 5 days on a growth medium. Explants were then transferred to liquid medium comprising 30 mg/L aspirin or acetylsalicylic acid (resulting in oxidative stress conditions) for one day. In control experiments, hypocotyls of non-transgenic *Brassica napus* plants N90-740 were cultivated on the same growth medium and then incubated for one day in liquid medium comprising 30 mg/L aspirin. In addition, hypocotyls of both the transgenic lines and the control line were cultivated on the same growth medium without aspirin.

[93] After the cultivation period, the ATP content of 125 explants was determined for each experiment. Additionally, the oxygen consumed in 3 hours by 125 explants was determined. The results are summarized in Table 1. The standard error of the mean was less than 6%. Whereas, the ratio of moles ATP per mg consumed oxygen in the control plants decreased in the control plants when oxidative stress was applied, the same ratio in the stress tolerant transgenic plant lines actually increased under stress conditions, and was considerably higher (about 24%) than in the control plants. The stress resistant transgenic lines thus maintained a constant energy production efficiency, whereas the control lines exhibited a decreased energy production efficiency. In addition, superoxide production, expressed as a percentage of superoxide production in control plants not subjected to the oxidative stress, did not increase in stress tolerant plants subjected to stress conditions.

[94] Table 1. Influence of stress on energy production efficiency of 5 days cultured *Brassica napus* hypocotyl explants.

Plant line	Stress	moles ATP per 125 explants	O ₂ mg/L consumed in 3 hrs by 125 explants	moles ATP	Superoxide production
				mg consumed O ₂	
N90-740 (control)	None	12.4×10^{-7}	2.96	4.19×10^{-7}	100%
	30mg/L aspirin	13.2×10^{-7}	4.06	3.25×10^{-7}	167%
Transgenic line	None	9.3×10^{-7}	2.33	3.99×10^{-7}	108%
	30mg/L aspirin	11.4×10^{-7}	2.82	4.04×10^{-7}	100%

[95] In another experiment, the NAD⁺ and ATP content of 4 different transgenic *Arabidopsis* lines comprising PARP gene expression reducing chimeric genes as described in WO 00/04173 were determined under high and low light conditions, and compared to the values obtained for a non transformed control line under the same conditions. The 4 different lines exhibited different degrees of stress resistance as exhibited e.g. by their ability to withstand heat and/or drought conditions. The values obtained for the NAD and ATP contents under high light stress are expressed as a percentage of the values for the NAD and ATP contents under low light conditions, and are plotted in Figure 2.

[96] The results show that high light stress leads to a significant NAD reduction in control plant cells and in the transgenic plant line which is the least stress resistant. The more stress resistant the transgenic plant lines are, the less significant the NAD reduction is under high light stress conditions.

[97] In another experiment, the NAD⁺ and ATP content of a segregating population resulting from a cross between transgenic corn lines comprising PARP gene expression reducing chimeric genes as described in WO 00/04173 and an untransformed corn line, were determined under conditions of nutrient (nitrogen) depletion, and compared to the values obtained for a non transformed control line under the same conditions. Figure 3 is a graphic representation of the of the obtained results. Hemizygous and azygous lines were discriminated by verification for the presence of the selectable marker gene. The NAD and ATP content was significantly higher in the hemizygous, stress tolerant plants than in the untransformed control plants or the azygous plants.

Example 2. Construction of ParG gene expression reducing chimeric genes.

[98] To reduce the expression of the PARG gene e.g. in *Arabidopsis* and related plants, a chimeric gene was constructed which is capable expressing a dsRNA comprising both a sense and antisense region which can form a double stranded RNA. Such dsRNA is very effective in reducing the expression of the genes with which is shares sequence homology, by post-transcriptional silencing. The chimeric gene comprises the following DNA fragments:

- A promoter region from Cauliflower mosaic Virus (CaMV 35S);
- A DNA fragment comprising 163 bp from the ParG gene from *Arabidopsis thaliana* in direct orientation (Genbank Accession number AF394690 from nucleotide position 973 to 1135);
- A DNA fragment encoding intron 2 from the pdk gene from *Flaveria*;
- The DNA fragment comprising 163 bp from the ParG gene from *Arabidopsis thaliana* in inverted orientation (Genbank Accession number AF394690 from nucleotide position 973 to 1135)
- A fragment of the 3' untranslated end from the octopine synthetase gene from *Agrobacterium tumefaciens*.

[99] This chimeric gene was introduced in a T-DNA vector, between the left and right border sequences from the T-DNA, together with a selectable marker gene providing resistance to the herbicide phosphinotricin.

[100] To reduce the expression of the PARG gene e.g. in potatoes and related plants, a chimeric gene is constructed which is capable expressing a dsRNA comprising both a sense and antisense region of a cDNA sequence from potato, that is capable of encoding a protein having high sequence identity with the N-terminal part of the *Arabidopsis* PARG protein. The chimeric gene comprises the following DNA fragments:

- A promoter region from Cauliflower mosaic Virus (CaMV 35S);
- A DNA fragment comprising a sequence of at least 100 bp from ParG homologue from *Solanum tuberosum* in direct orientation (Genbank Accession number BE340510);
- A DNA fragment encoding intron 2 from the pdk gene from *Flaveria*;
- The DNA fragment comprising the sequence of at least 100 bp from ParG homologue from *Solanum tuberosum* in inverted orientation (Genbank Accession number BE340510);
- A fragment of the 3' untranslated end from the octopine synthetase gene from *Agrobacterium tumefaciens*

[101] This chimeric gene is introduced in a T-DNA vector, between the left and right border sequences from the T-DNA, together with a selectable marker gene providing resistance to the herbicide phosphinotricin.

Example 3. Analysis of transgenic plant lines comprising ParG gene expression reducing chimeric genes.

[102] The chimeric genes of Example 2 are introduced into *Arabidopsis* or potato respectively, by *Agrobacterium* mediated transformation.

[103] The population of obtained transgenic lines is subjected to the following stress conditions, together with control plants:

- Increased heat for a period of days (greenhouse) or hours (in vitro)
- Drought for a period of days
- High light conditions for a period of days
- Nutrient depletion

[104] Individual plant lines surviving well the above mentioned stress conditions are selected.

[105] The NAD content and ATP content for the above mentioned plants is determined under control and stress conditions.

Example 4. Quantitative determination of NAD, ATP and superoxide radicals in plant cells.

[106] Quantification of ATP in plant tissues was done basically as described by Rawlyer et al. (1999), Plant Physiol. 120, 293-300. The assay was used for the determination of the ATP content of hypocotyl explants that were cultured for 4-5 days on A2S3 medium or 2 weeks old *in vitro* cultured *Arabidopsis* plants. All manipulations are performed on crushed ice unless otherwise indicated.

[107] ATP extraction

- Freeze plant material with liquid nitrogen
 - 100 hypocotyl explants
 - \pm 700mg *Arabidopsis* plants (roots + shoots) (about 32-37 18-days old C24 plants)
- Put frozen hypocotyls in mortar and add 6ml of 6% perchloric acid.
- Extraction can be done at room temperature using a pestle. After extraction, put samples as soon as possible on ice.
- Centrifuge at 24,000g (Sorvall, SS34 rotor at 14,000rpm) for 10min. at

4°C.

- The supernatant is neutralized with 5M K₂CO₃ (add 350µl of 5M K₂CO₃ to 3ml of supernatant).
- KClO₄ is removed by spinning as described above.

[108] Quantitative bioluminescent determination of ATP

- The ATP bioluminescent assay kit from Sigma is used (FL-AA).
- Dilute extract 6000 x (about 6 mL extract from which 100µl is taken, that is diluted 1000 times) The dilutions are made with the 'ATP assay mix dilution buffer' (FL-AAB) of the ATP bioluminescent assay kit
- The amount of light that is produced is measured with the TD-20/20 luminometer of Turner Designs (Sunnyvale, USA).
- Standard curve: dissolve ATP standard of kit (FL-AAS) in 10ml of water (2x10⁻⁶ moles)

[109] Quantification of NAD⁺ and NADH in plant tissues was performed, essentially as described by Karp et al. (1983) or Filipovic et al. (1999) on the following plant material:

Brassica napus: 150 5-days cultured hypocotyl explants/sample

Arabidopsis: 1000mg 18-days old in vitro grown plants (shoots + roots)/sample (corresponds to ±60 C24 plants)

[110] Assay solution

(A) For measuring NADH: 25mM potassium phosphate buffer pH7

0.1mM DTT

3µM FMN (Fluka, 83810)

30µM n-decanal (Sigma, D-7384)

(B) For measuring NAD⁺ + NADH:

idem as for measuring NADH alone

+ 2µg/mL alcohol dehydrogenase (Roche, 102 717)

[111] Extraction

- Freeze with liquid nitrogen
- Put frozen plant material in cooled mortar (cooled at -20°C) and add 5mL extraction buffer
- Grind material using a pestle
- Centrifuge at 24 000g (Sorvall, SS34 rotor at 14 000rpm) for 15 minutes at 4°C
- Take 1 mL of supernatant for analysis

[112] Assay***NADH***

- 390 μL of assay solution A
- + 10 μL extract
- + 2 μL NAD(P)H:FMN oxidoreductase
- + 100 μL luciferase solution

NAD⁺ + NADH

- 390 μL of assay solution B
- + 10 μL extract
- 2 minutes at room temperature
- + 2 μL NAD(P)H:FMN oxidoreductase
- + 100 μL luciferase solution

The amount of light that is produced is measured with the TD-20/20 luminometer of Turner Designs (Sunnyvale, USA)

NADH-standard

NADH stock solution: 1mM (7.1mg/10mL H_2O)

NADH: disodium salt, Roche, 107 735

Dilution series in 10mM potassium phosphate buffer pH7: (10^{-2}); 5×10^{-3} ; 2×10^{-3} ; 10^{-3} ; 5×10^{-4}

Add 10 μL of dilutions in 390 μL of assay solution A and perform reaction

Make standard curve

- [113]** Superoxide radicals production was measured by quantifying the reduction of XTT as described in De Block and De Brouwer (2002) Plant Physiol. Biochem. 40, 845-852

[114] BRASSICA NAPUS**[115] Media and reaction buffers****Sowing medium (medium 201):**

Half concentrated Murashige and Skoog salts

2% sucrose

pH 5.8

0.6% agar (Difco Bacto Agar)

250mg/l triacillin

Callus inducing medium A2S3:

MS medium, 0.5g/l Mes (pH 5.8), 3% sucrose, 40mg/l adenine-SO₄,

0.5% agarose, 1mg/l 2,4-D, 0.25mg/l NAA, 1mg/l BAP, 250mg/l

triacillin

Incubation medium:

25mM K-phosphate buffer pH5.8

2% sucrose

1 drop Tween20 for 25ml medium

Reaction buffer:

50mM K-phosphate buffer pH7.4

1mM sodium,3'-{1-[phenylamino-carbonyl]-3,4-tetrazolium}-bis(4-methoxy-6-nitro) = XTT (bts, Germany, cat n° 2525)

1 drop Tween20 for 25ml buffer

[116] Sterilization of seeds - pregermination of seeds - growing of the seedlings. Seeds are soaked in 70% ethanol for 2 min, then surface-sterilized for 15 min in a sodium hypochlorite solution (with about 6% active chlorine) containing 0.1% Tween20. Finally, the seeds are rinsed with 1l of sterile tap water. Incubate seeds for at least one hour in sterile tap water (to allow diffusion from seeds of components that may inhibit germination). Seeds are put in 250ml erlenmeyer flasks containing 50ml of sterile tap water (+ 250mg/l triacillin). Shake for

about 20 hours. Seeds from which the radicle is protruded are put in Vitro Vent containers from Duchefa containing about 125ml of sowing medium (10 seeds/vessel, not too many to reduce loss of seed by contamination). The seeds are germinated at $\pm 24^{\circ}\text{C}$ and 10-30 Einstein/s⁻¹m⁻² with a daylength of 16h.

[117] Preculture of the hypocotyl explants and induction of stress

- 12-14 days after sowing, the hypocotyls are cut in about 7-10mm segments.
- The hypocotyl explants (25 hypocotyls/Optilux Petridish, Falcon S1005, Denmark) are cultured for 5 days on medium A2S3 at 25°C (at 10-30 Einstein/s⁻¹m⁻²).

[118] XTT-assay

- Transfer 150 hypocotyl explants to a 50ml Falcon tube.
- Wash with reaction buffer (without XTT).
- Add 20mL reaction buffer + XTT.
(explants have to be submerged, but do not vacuum infiltrate)
- Incubate in the dark at 26°C for about 3hours
- Measure the absorption of the reaction medium at 470nm

[119] *ARABIDOPSIS THALIANA*

[120] Media and reaction buffers

Plant medium:

Half concentrated Murashige and Skoog salts

B5 vitamins

1.5% sucrose

pH 5.8

0.7% Difco agar

Incubation medium:

10mM K-phosphate buffer pH5.8

2% sucrose

1 drop Tween20 for 25ml medium

Reaction buffer:

50mM K-phosphate buffer pH7.4

1mM sodium,3'-[1-[phenylamino-carbonyl]-3,4-tetrazolium]-bis(4-methoxy-6-nitro) = XTT (bts, Germany, cat n° 2525)

1 drop Tween20 for 25ml buffer

[121] *Arabidopsis* plants

- *Arabidopsis* lines: control

lines to test

- Sterilization of *Arabidopsis* seeds:

2min. 70% ethanol

10 min. bleach (6% active chlorine) + 1drop Tween 20 for 20ml solution

wash 5 times with sterile tap water

- Pregermination of seeds:

In 9cm Optilux Petridishes (Falcon) containing 12ml sterile tap water.

Low light overnight to 24 hours.

- Growing of *Arabidopsis* plants

Seeds are sown in Intergrid Tissue Culture disks of Falcon (nr. 3025) containing ± 125 ml of plant medium: 1 seed/grid.

Plants are grown at 24°C

$30\mu\text{Einstein s}^{-1}\text{m}^{-2}$

16hours light - 8hours dark

for about 3 weeks (before bolting)

[122] XTT-assay

Control condition (no stress)

- Harvest shoots (roots included) from agar plates and put them directly in a 50ml Falcon tube containing reaction buffer (without XTT)

Stressed shoots

- Transfer shoots to 50ml Falcon tubes containing reaction buffer (without XTT)

- Replace reaction buffer with buffer containing XTT (40mL/tube)

- Shoots have to be submerged, but do not vacuum infiltrate
- Incubate in the dark at 26°C for about 3hours
- Measure the absorption of the reaction medium at 470nm

[123] Quantification of respiration by measuring oxygen consumption using a Clark polarographic electrode was done in the following way:

[124] Plant material

Brassica napus

150-200* hypocotyl explants

Cultured for 5 days at 25°C

(*cfr.* protocol vigour assay)

* 150 explants error <10%; 200 explants error <6%

Arabidopsis

For $C24 \pm 1000\text{mg}^*$ *in vitro* plants (shoots + roots) (corresponds with ~50 18-days old plants)

Pregerminate seeds before sowing

Grow for 18 days at 24°C

(*cfr.* protocol *in vitro* growth Arabidopsis)

* for error <8%

[125] Incubation media

Brassica napus

25mM K-phosphate buffer pH5.8

2% sucrose

Tween20 (1 drop/25ml)

Arabidopsis

10mM K-phosphate buffer pH5.8

2% sucrose

$\mu\text{mol m}^{-1} \text{s}^{-1}$). After 1 day, the NAD and NADH contents were determined using the enzymatic cycling method (Karp et al. (1983) Anal. Biochem. 128, pp 175-180). A portion of the seedlings were cultivated further under high light conditions for about 3 to about days, after which the damage was scored. Damage was visible as darkening of the young leaves and shoot tip, bleaching of older leaves and growth retardation. The results are summarized in Table 1 for Arabidopsis and in Table 2 for tobacco.

[129] Table 1. Analysis of Arabidopsis (Columbia). $\pm R$ indicates that some dark pigmentation was observed. ND: not determined

	High light tolerance	NAD+NADH content in 1 gram of tissue ($10^{-3}\mu\text{M}$)	% TTC-reducing capacity vs control
Non-transgenic control	S	17.3	100
Transgenic line 9	R	28.2	ND
Transgenic line 10	R	31.7	ND
Transgenic line 11	$\pm R$	26.5	ND
Transgenic line 12	S	19.4	ND
Transgenic line 26	R	33.2	55
Transgenic line 27	S	21.3	100
Transgenic line 28	$\pm R$	26.5	75
Transgenic line 29	S	17.7	102
Transgenic line 30	R	28.3	66

[130] Table 2. Analysis of *Nicotiana tabacum* c.v. Petit Havana SR1. \pm R indicates that some dark pigmentation was observed. R/S indicates that the resistance phenotype was not very clear.

	High light tolerance	% TTC-reducing capacity vs control
Non-transgenic control	S	100
Transgenic line 1	R/S	88
Transgenic line 2	\pm R	79
Transgenic line 3	R	53

[131] There is a positive correlation between the resistance to high light stress in the transgenic plants and the NAD⁺/NADH content of the cells. An inverse correlation can be seen between TTC reducing capacity and high light tolerance.

Example 6. Construction of ParG gene expression reducing chimeric genes suited for use in cereal plants.

[132] To reduce the expression of the PARG gene e.g. in cereals such as rice or corn (maize) and related plants, a chimeric gene is constructed which is capable expressing a dsRNA comprising both a sense and antisense region of nucleotide sequence from rice, that is capable of encoding a protein having high sequence identity with PARG protein encoding nucleotide sequences. The chimeric gene comprises the following DNA fragments:

- A promoter region from Cauliflower mosaic Virus (CaMV 35S);
- A DNA fragment comprising a sequence of at least 100 bp from ParG homologue from *Oryza sativa* (SEQ ID No 15) in direct orientation;
- A DNA fragment encoding intron 2 from the pdk gene from *Flaveria*;

- A DNA fragment comprising a sequence of at least 100 bp from ParG homologue from *Oryza sativa* (SEQ ID No 15) in inverted orientation;
- A fragment of the 3' untranslated end from the octopine synthetase gene from *Agrobacterium tumefaciens*.

[133] This chimeric gene is introduced in a T-DNA vector, between the left and right border sequences from the T-DNA, together with a selectable marker gene providing resistance to e.g. the herbicide phosphinotricin.

[134] To reduce the expression of the PARG gene e.g. in cereals such as rice or corn (maize) and related plants, a chimeric gene is constructed which is capable expressing a dsRNA comprising both a sense and antisense region of nucleotide sequence from rice, that is capable of encoding a protein having high sequence identity with PARG protein encoding nucleotide sequences. The chimeric gene comprises the following DNA fragments:

- A promoter region from Cauliflower mosaic Virus (CaMV 35S);
- A DNA fragment comprising a sequence of at least 100 bp from ParG homologue from *Zea mays* (SEQ ID No 23) in direct orientation;
- A DNA fragment encoding intron 2 from the pdk gene from *Flaveria*;
- A DNA fragment comprising a sequence of at least 100 bp from ParG homologue from *Zea mays* (SEQ ID No 23) in inverted orientation;
- A fragment of the 3' untranslated end from the octopine synthetase gene from *Agrobacterium tumefaciens*.

[135] This chimeric gene is introduced in a T-DNA vector, between the left and right border sequences from the T-DNA, together with a selectable marker gene providing resistance to e.g. the herbicide phosphinotricin. The nucleotide sequence of two examples of such T-DNA vectors

comprising two different chimeric genes as described in the previous paragraph is represented in SEQ ID Nos 24 and 25.

Example 7. Analysis of transgenic plant lines comprising ParG gene expression reducing chimeric genes.

[136] The chimeric genes of Example 6 are introduced into rice or corn respectively, by Agrobacterium mediated transformation.

[137] The population of obtained transgenic lines is subjected to the following stress conditions, together with control plants:

- Increased heat for a period of days (greenhouse) or hours (in vitro)
- Drought for a period of days
- High light conditions for a period of days
- Nutrient depletion

[138] Individual plant lines surviving well the above mentioned stress conditions, or at least one thereof, are selected.

[139] The NAD content and ATP content for the above mentioned plants is determined under control and stress conditions.

Claims:

1. A method to produce a plant tolerant to stress conditions comprising the steps of

(a) providing plant cells with a chimeric gene to create transgenic plant cells, said chimeric gene comprising the following operably linked DNA fragments

- (i) a plant-expressible promoter;
- (ii) a DNA region, which when transcribed yields an ParG inhibitory RNA molecule;
- (iii) a 3' end region involved in transcription termination and polyadenylation;

(b) regenerating a population of transgenic plant lines from said transgenic plant cell; and

(c) identifying a stress tolerant plant line within said population of transgenic plant lines.

2. The method according to claim 1, wherein said parG inhibitory RNA molecule comprises comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present in said plant cell.

3. The method according to claim 1, wherein said parG inhibitory RNA molecule is comprises comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell.

4. The method according to claim 2 or 3, wherein said chimeric gene further comprises a DNA region encoding a self-splicing ribozyme between said DNA region coding for said parG inhibitory RNA molecule and said 3' end region.

5. The method according to claim 1, wherein said parG inhibitory RNA comprises a sense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the nucleotide sequence of the ParG gene present

in said plant cell and an antisense region comprising a nucleotide sequence of at least 20 consecutive nucleotides of the complement of the nucleotide sequence of the ParG gene present in said plant cell, wherein said sense and antisense region are capable of forming a double stranded RNA region comprising said at least 20 consecutive nucleotides.

6. The method according to any one of claims 1 to 5 wherein said stress conditions is selected from heat, drought, nutrient depletion, oxidative stress or high light conditions.

7. The method according to any one of claims 1 to 6, comprising further crossing said transgenic plant line with another plant line to obtain stress tolerant progeny plants.

8. A method to produce a plant tolerant to stress conditions comprising the steps of:

(a) isolating a DNA fragment of at least 100 bp comprising a part of the parG encoding gene of said plant;

(b) producing a chimeric gene by operably linking the following DNA fragments;

(i) a plant expressible promoter region;

(ii) said isolated DNA fragment comprising part of the parG encoding gene of said plant in direct orientation compared to the promoter region;

(iii) said isolated DNA fragment comprising part of the parG encoding gene of said plant in inverted orientation compared to the promoter region;

(iv) a 3' end region involved in transcription termination and polyadenylation;

(c) providing plant cells with said chimeric gene to create transgenic plant cells

(d) regenerating a population of transgenic plant lines from said transgenic plant cell; and

(e) identifying a stress tolerant plant line within said population of transgenic plant lines.

9. A DNA molecule comprising

- (i) a plant-expressible promoter;
- (ii) a DNA region, which when transcribed yields a ParG inhibitory RNA molecule;
- (iii) a 3' end region involved in transcription termination and polyadenylation.

10. The DNA molecule according to claim 9, wherein said DNA region comprises a nucleotide sequence of at least 21 to 100 nucleotides of a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID No 1, 2 or 16 or at least 21 to 100 nucleotides of a nucleotide sequence of SEQ ID 3, 4, 15 or 23.

11. A plant cell comprising the DNA molecule of claim 9 or 10.

12. A plant consisting essentially of the plant cells of claim 11.

13. A process for producing stress tolerant plants, comprising the step of further crossing a plant of claim 12 with another plant.

14. Seeds and propagating material of a plant according to claim 12, comprising the chimeric gene of claim 9 or 10.

15. Plants obtainable or obtained by the process of claim 8.

16. A method to produce a plant tolerant to stress conditions comprising the steps of

(a) providing plant cells with a chimeric gene to create transgenic plant cells, said chimeric gene comprising the following operably linked DNA fragments

- (i) a plant-expressible promoter;

(ii) a DNA region, which when transcribed yields an ParG inhibitory RNA molecule, said DNA region comprising a nucleotide sequence of at least 21 to 100 nucleotides of a nucleotide sequence encoding a protein comprising the amino acid sequence of SEQ ID No 1, 2 or 16 or at least 21 to 100 nucleotides of a nucleotide sequence of SEQ ID 3, 4, 15 or 23;

(iii) a 3' end region involved in transcription termination and polyadenylation;

(b) regenerating a population of transgenic plant lines from said transgenic plant cell; and

(c) identifying a stress tolerant plant line within said population of transgenic plant lines.

17. A method to produce a plant tolerant to stress conditions comprising the steps of

(a) subjecting a plant cell line or a plant or plant line, to mutagenesis;

(b) identifying those plant cells or plants that have a mutation in an endogenous ParG gene;

(c) subjecting the identified plant cells or plants to stress conditions;

(d) identifying plant cells or plants that tolerate said stress conditions better than control plants.

18. A method to produce a plant tolerant to stress conditions comprising the steps of

(a) selecting a plant cell line or a plant or plant line which is resistant to a ParG inhibitor;

(b) identifying those plant cells or plants that have a mutation in an endogenous ParG gene;

(c) subjecting the identified plant cells or plants to stress conditions;

(d) identifying plant cells or plants that tolerate said stress conditions better than control plants.

19. A stress tolerant plant cell or plant comprising a mutation in an endogenous ParG gene.





Figure 1






Figure 3

SEQUENCE LISTING

J05 Rec'd PCT/PTO 07 OCT 2005

<110> Bayer BioScience N.V.
De Block, Marc

<120> Methods and means for increasing the tolerance of plants to stress conditions.

<130> BCS 03 2002 WO1

<150> EP03076044.1

<151> 2003-04-09

<150> US 60/496,688

<151> 2003-08-21

<160> 25

<170> PatentIn version 3.1

<210> 1

<211> 548

<212> PRT

<213> Arabidopsis thaliana

<400> 1

```

Met Glu Asn Arg Glu Asp Leu Asn Ser Ile Leu Pro Tyr Leu Pro Leu
1      5      10      15
Val Ile Arg Ser Ser Leu Tyr Trp Pro Pro Arg Val Val Glu Ala
      20      25      30
Leu Lys Ala Met Ser Glu Gly Pro Ser His Ser Gln Val Asp Ser Gly
      35      40      45
Glu Val Leu Arg Gln Ala Ile Phe Asp Met Arg Arg Ser Leu Ser Phe
      50      55      60
Ser Thr Leu Glu Pro Ser Ala Ser Asn Gly Tyr Ala Phe Leu Phe Asp
      65      70      75      80
Glu Leu Ile Asp Glu Lys Glu Ser Lys Arg Trp Phe Asp Glu Ile Ile
      85      90      95
Pro Ala Leu Ala Ser Leu Leu Leu Gln Phe Pro Ser Leu Leu Glu Val
      100     105     110
His Phe Gln Asn Ala Asp Asn Ile Val Ser Gly Ile Lys Thr Gly Leu
      115     120     125
Arg Leu Leu Asn Ser Gln Gln Ala Gly Ile Val Phe Leu Ser Gln Glu
      130     135     140
Leu Ile Gly Ala Leu Leu Ala Cys Ser Phe Phe Cys Leu Phe Pro Asp
      145     150     155     160
Asp Asn Arg Gly Ala Lys His Leu Pro Val Ile Asn Phe Asp His Leu
      165     170     175
Phe Ala Ser Leu Tyr Ile Ser Tyr Ser Gln Ser Gln Glu Ser Lys Ile
      180     185     190
Arg Cys Ile Met His Tyr Phe Glu Arg Phe Cys Ser Cys Val Pro Ile
      195     200     205
Gly Ile Val Ser Phe Glu Arg Lys Ile Thr Ala Ala Pro Asp Ala Asp
      210     215     220
Phe Trp Ser Lys Ser Asp Val Ser Leu Cys Ala Phe Lys Val His Ser
      225     230     235     240
Phe Gly Leu Ile Glu Asp Gln Pro Asp Asn Ala Leu Glu Val Asp Phe
      245     250     255
Ala Asn Lys Tyr Leu Gly Gly Gly Ser Leu Ser Arg Gly Cys Val Gln
      260     265     270
Glu Glu Ile Arg Phe Met Ile Asn Pro Glu Leu Ile Ala Gly Met Leu

```

275 280 285
 Phe Leu Pro Arg Met Asp Asp Asn Glu Ala Ile Glu Ile Val Gly Ala
 290 295 300
 Glu Arg Phe Ser Cys Tyr Thr Gly Tyr Ala Ser Ser Phe Arg Phe Ala
 305 310 315 320
 Gly Glu Tyr Ile Asp Lys Lys Ala Met Asp Pro Phe Lys Arg Arg Arg
 325 330 335
 Thr Arg Ile Val Ala Ile Asp Ala Leu Cys Thr Pro Lys Met Arg His
 340 345 350
 Phe Lys Asp Ile Cys Leu Leu Arg Glu Ile Asn Lys Ala Leu Cys Gly
 355 360 365
 Phe Leu Asn Cys Ser Lys Ala Trp Glu His Gln Asn Ile Phe Met Asp
 370 375 380
 Glu Gly Asp Asn Glu Ile Gln Leu Val Arg Asn Gly Arg Asp Ser Gly
 385 390 395 400
 Leu Leu Arg Thr Glu Thr Thr Ala Ser His Arg Thr Pro Leu Asn Asp
 405 410 415
 Val Glu Met Asn Arg Glu Lys Pro Ala Asn Asn Leu Ile Arg Asp Phe
 420 425 430
 Tyr Val Glu Gly Val Asp Asn Glu Asp His Glu Asp Asp Gly Val Ala
 435 440 445
 Thr Gly Asn Trp Gly Cys Gly Val Phe Gly Gly Asp Pro Glu Leu Lys
 450 455 460
 Ala Thr Ile Gln Trp Leu Ala Ala Ser Gln Thr Arg Arg Pro Phe Ile
 465 470 475 480
 Ser Tyr Tyr Thr Phe Gly Val Glu Ala Leu Arg Asn Leu Asp Gln Val
 485 490 495
 Thr Lys Trp Ile Leu Ser His Lys Trp Thr Val Gly Asp Leu Trp Asn
 500 505 510
 Met Met Leu Glu Tyr Ser Ala Gln Arg Leu Tyr Lys Gln Thr Ser Val
 515 520 525
 Gly Phe Phe Ser Trp Leu Leu Pro Ser Leu Ala Thr Thr Asn Lys Ala
 530 535 540
 Ile Gln Pro Pro
 545
 <210> 2
 <211> 169
 <212> PRT
 <213> Solanum tuberosum

<400> 2
 Met Glu Asn Arg Glu Asp Val Lys Ser Ile Leu Pro Phe Leu Pro Val
 1 5 10 15
 Cys Leu Arg Ser Ser Ser Leu Phe Trp Pro Pro Leu Val Val Glu Ala
 20 25 30
 Leu Lys Ala Leu Ser Glu Gly Pro His Tyr Ser Asn Val Asn Ser Gly
 35 40 45
 Gln Val Leu Phe Leu Ala Ile Ser Asp Ile Arg Asn Ser Leu Ser Leu
 50 55 60
 Pro Asp Ser Ser Ile Ser Ser Ser Ala Ser Asp Gly Phe Ser Leu Leu
 65 70 75 80
 Phe Asp Asp Leu Ile Pro Arg Asp Glu Ala Val Lys Trp Phe Lys Glu
 85 90 95
 Val Val Pro Lys Met Ala Asp Leu Leu Leu Arg Leu Pro Ser Leu Leu
 100 105 110
 Glu Ala His Tyr Glu Lys Ala Asp Gly Gly Ile Val Lys Gly Val Asn
 115 120 125

Thr Gly Leu Arg Leu Leu Glu Ser Gln Gln Pro Gly Ile Val Phe Leu
 130 135 140

Ser Gln Glu Leu Val Gly Ala Leu Leu Ala Cys Ser Phe Phe Cys Tyr
 145 150 155 160

Ser Leu Pro Met Ile Glu Val Ser Val
 165

<210> 3
 <211> 1647
 <212> DNA
 <213> Arabidopsis thaliana

<400> 3
 atggagaatc gcgaagatct taactcaatt cttccgtacc ttccacttgt aattcgttcg 60
 tcgctcgctgt attggccgcc gcgtgtggtg gaggcggttaa aggcaatgtc tgaaggacca 120
 tctcacagcc aagttgactc aggagaggtt ctacggcaag ctattttcga tatgagacga 180
 tccttatctt tctctactct cgagccatct gcttctaatt gctacgcatt tctctttgac 240
 gaattgattg atgagaaaga atcaaagaga tggttcgatg agattatccc agcattggcg 300
 agcttacttc tacagtttcc atctctgtta gaagtgcatt tccaaaatgc tgataatatt 360
 gttagtggaa tcaaaaccgg tcttcgtttg ttaaattccc aacaagctgg cattgttttc 420
 ctacgccagg agttgattgg agctcttctt gcattgctct tcttttggtt gtttcgggat 480
 gataatagag gtgcaaaaca ccttcagtc atcaactttg atcatttgtt tgcaagcctt 540
 tatataagtt atagtcaaag tcaagaaagc aagataagat gtattatgca ttactttgaa 600
 aggttttgct cctgcgtgcc tattggtatt gtttcatttg aacgcaagat taccgctgct 660
 cctgatgctg atttctggag caagtctgac gtttctcttt gtgcatttaa ggttcaactc 720
 tttgggttaa ttgaagatca acctgacaat gctctcgaag tggactttgc aaacaagtat 780
 ctcgagggtg gttccctaag tagagggtgc gtgcaggaag agatacgtt catgattaac 840
 cctgaattaa tcgctggcat gcttttcttg cctcggatgg atgacaatga agctatagaa 900
 atagtgtgtg cggaaagatt ttcatgttac acagggtatg catcttcggt tcggtttgct 960
 ggtgagtaca ttgacaaaaa ggcaatggat cctttcaaaa ggcaagaac cagaattgtt 1020
 gcaattgatg cattatgtac accgaagatg agacacttta aagatatatg tcttttaagg 1080
 gaaattaata aggcactatg tggcttttta aattgttaga aggcttggga gcaccagaat 1140
 atattcatgg atgaaggaga taatgaaatt cagcttgtcc gaaacggcag agattctggt 1200
 cttctgcgta cagaaactac tgcgtcacac cgaactccac taaatgatgt tgagatgaat 1260
 agagaaaagc ctgctaacaa tcttatcaga gattttatg tggaggagt tgataacgag 1320
 gatcatgaag atgatggtgt cgcgacaggg aattggggat gtggtgtttt tggaggagac 1380
 ccagagctaa aggctacgat acaatggctt gctgcttccc agactcgaag accatttata 1440
 tcatattaca cctttggagt agaggcactc cgaaacctag atcagggtgac gaagtggatt 1500
 ctttcccata aatggactgt tggagatctg tggaaatga tgtagaata ttctgctcaa 1560
 aggtctaca agcaaaccag tggtggcttc ttttcttggc tacttccatc tctaagctacc 1620
 accaacaag ctatccagcc gccttga 1647

<210> 4
 <211> 598
 <212> DNA
 <213> Solanum tuberosum

<400> 4
 gcaatggaga atagagaaga cgtgaagtca atccttccct ttttgccggt gtgtctccga 60
 tcatcttctc ttttctggcc gccgctagtt gttgaagcac tgaaagccct ctctgaaggc 120
 cctcattaca gcaatgttaa ctccggccaa gtccctcttc tcgcaatctc cgacattcgg 180
 aattcccttt cactacctga ttcttcaatt tctcttctg cttcagacgg attttctctc 240
 ttattttgat atttaattcc tagggatgaa gctgttaaat ggttcaaaga agtgggtgccg 300
 aaaaatggcg atttgctatt gcgggtgcct tcttatttgg aggtcacta tgagaaggct 360
 gatgggtgaa ttgttaaagg agtcaacact ggtcttcgct tatttggaac acaacagcct 420
 ggcattgttt tctcagtcga ggaattagtc ggtgctcttc ttgcatgttc cttcttttgc 480
 tattccctac caatgataga ggtatctgta tgatcagtat gacgagaaat ttgaaaataa 540
 attgaagtgc attcttcaat attttgagag gattggctca ttgatacctg cgggctac 598

<210> 5
 <211> 37
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ParGAt1
 <400> 5
 ggatccctg caggacaaaa aggcaatgga tccttttc

37

<210> 6
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ParGAt2
 <400> 6
 gcacgaattc ggggccgcg tgctcccaag ccttgctac

39

<210> 7
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ParGSt1
 <400> 7
 ggatccctg caggctcact atgagaaggc tgatgggtgg

39

<210> 8
 <211> 43
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> oligonucleotide primer ParGSt2
 <400> 8
 gcacgaattc ggggccgct catactgatac atacagatac etc

43

<210> 9
 <211> 13466
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> nucleotide sequence of pTVE428
 <220>
 <221> misc_feature
 <222> (198)..(222)
 <223> Right T-DNA border

<220>
 <221> misc_feature
 <222> (983)..(273)
 <223> 3' ocs (3' untranslated end of octopine synthase gene)

<220>
 <221> misc_feature
 <222> (995)..(1155)
 <223> part of poly (ADP-ribose) glycohydrolase

<220>
 <221> misc_feature
 <222> (1929)..(1188)
 <223> intron 2 from the Pdk gene of Flaveria

 <220>
 <221> misc_feature
 <222> (2122)..(1962)
 <223> part of poly (ADP-ribose) glycohydrolase

 <220>
 <221> misc_feature
 <222> (3476)..(2131)
 <223> 35S promoter region from Cauliflower Mosaic Virus

 <220>
 <221> misc_feature
 <222> (3948)..(3737)
 <223> 3' untranslated end of gene 7 from Agrobacterium tumefaciens

 <220>
 <221> misc_feature
 <222> (4521)..(3970)
 <223> bar coding region

 <220>
 <221> misc_feature
 <222> (6247)..(4522)
 <223> PSSuAra promoter region

 <220>
 <221> misc_feature
 <222> (6415)..(6439)
 <223> left border region of T-DNA of Agrobacterium tumefaciens

<400> 9
 agattcgaag ctccggtcccg tgggtgttct gtcgtctcgt tgtacaacga aatccattcc 60
 cattccgcgc tcaagatggc ttcccctcgg cagttcatca gggctaaatc aatctagccg 120
 acttgtcccg tgaaatgggc tgcactccaa cagaaacaat caaacaacaa tacacagcga 180
 cttattcaca cgcgacaaat tacaacggta tatatcctgc cagtactcgg ccgtcgcaccg 240
 cggtagcccg gaattaagct tgcattgcctg caggtcctgc tgagcctcga catgttgtcg 300
 caaaattcgc cctggaccgg cccaacgatt tgcgtcact gtcaagggtt gacctgcact 360
 tcatttgggg ccacataca ccaaaaaaat gctgcataat tctcggggca gcaagtcggt 420
 taccgcggcg ccgtgctgga cggggttgaa tgggtgcccg aactttcggg agagcgggacg 480
 gccaatatc aacttcaagg aatctcacc atgcgcgcgg gcggggaacc ggagttccct 540
 tcagtgaacg ttattagttc gccgctcggg gtgtcgtaga tactagcccc tggggccctt 600
 tgaaatttga ataagattta tgtaatcagt cttttaggtt tgaccgggtc tgccgctttt 660
 tttaaaattg gatttgtaat aataaaacgc aattgtttgt tattgtggcg ctctatcata 720
 gatgtcgcta taaacctatt cagcacaata tattgttttc attttaatat tgtacatata 780
 agtagtaggg tacaatcagt aaattgaacg gagaatatta ttcataaaaa tacgatagta 840
 acgggtgata tattcattag aatgaaccga aaccggcggg aaggatctga gctacacatg 900
 ctcagggttt ttacaacgtg cacaacagaa ttgaaagcaa atatcatgcg atcataggcg 960
 tctcgcatat ctcattaaag caggactcta gagacaaaaa ggcaatggat cttttcaaaa 1020
 ggccaagaac cagaattggt gcaattgatg cattatgtac accgaagatg agacacttta 1080
 aagatatatg tcttttaagg gaaattaata aggcactatg tggcttttta aattgtagca 1140
 aggcttggga gcaccatcga ttccgaaccc agcttcccaa ctgtaataca tccaaatgta 1200
 agatcaatga taacacaatg acatgatcta tcatgttacc ttgtttattc atgttcgact 1260
 aattcattta attaatagtc aatccattta gaagttaata aaactacaag tattatttag 1320
 aaattaataa gaatgttgat tgaaaataat actatataaa atgatagatc ttgcgctttg 1380
 ttatattagc attagattat gttttgttac attagattac tgtttctatt agtttgatat 1440

tatttgttac	tttagcttgt	tatttaatat	tttgtttatt	gataaattac	aagcagattg	1500
gaattttctaa	caaaatgttt	attaactttt	aaactaaaat	atttagtaat	ggtatagata	1560
tttaattata	taataaacta	ttaatcataa	aaaaatatta	ttttaattta	tttattctta	1620
tttttactat	agtattttat	cattgatatt	taattcatca	aaccagctag	aattactatt	1680
atgattaaaa	caaataattaa	tgctagtata	tcatcttaca	tgttcgatca	aattcattaa	1740
aaataatata	cttactctca	acttttatct	tcttcgtctt	acacatcact	tgcatatttt	1800
ttttacatta	ctatgttggt	tatgtaaaca	atatatttat	aaattatttt	ttcacaatta	1860
taacaactat	attattataa	tcatacta	taacatcact	taactatttt	atactaaaag	1920
gaaaaaagaa	aataattatt	tccttaccaa	gctggggtag	cgggtgctcc	aagccttgct	1980
acaattttaa	aagccacata	gtgccttatt	aatttccctt	aaaagacata	tatctttaa	2040
gtgtctcatc	ttcggtgtac	ataatgcac	aattgcaaca	attctgggtc	ttcgctttt	2100
gaaaggatcc	attgcctttt	tgctctcgag	cgtgtcctct	ccaaatgaaa	tgaacttcc	2160
tatatagagg	aagggtcctg	cgaaggatag	tgggattgtg	cgtcatccct	tacgtcagtg	2220
gagatgtcac	atcaatccac	ttgctttgaa	gacgtggttg	gaacgtcttc	tttttccacg	2280
atgctcctcg	tgggtggggg	tccatctttg	ggaccactgt	cggcagagag	atcttgaatg	2340
atagcctttc	ctttatcgca	atgatggcat	ttgtaggagc	caccttccct	ttctactgtc	2400
ctttcgatga	agtgaacagat	agctgggcaa	tggaaatccga	ggaggtttcc	cgaattatc	2460
ctttgttgaa	aagtctcaat	agcccttttg	tcttctgaga	ctgtatcttt	gacatttttg	2520
gagtagacca	gagtgctgtg	ctccaccatg	ttgacgaaga	ttttctctct	gtcattgagt	2580
cgtaaaagac	tctgtatgaa	ctgttcgcca	gtcttcacgg	cgagttctgt	tagatcctcg	2640
atgtgaatct	tagactccat	gcattggcctt	agattcagta	ggaactacct	tttagagac	2700
tccaatctct	attacttgcc	ttggtttatg	aagcaagcct	tgaatcgctc	atactggaat	2760
agtacttctg	atcttgagaa	atatgtcttt	ctctgtgttc	ttgatgcaat	tagtctgaa	2820
tcttttgact	gcattcttaa	ccttcttggg	aaggattttg	atctcctgga	gattgttact	2880
cgggtagatc	gtcttgatga	gacctgtctg	gtaggcctct	ctaaccatct	gtgggtcagc	2940
attctttctg	aaattgaaga	ggctaaccct	ctcattatca	gtgggtgaaca	tagtgtcgtc	3000
accttcacct	tcgaacttcc	ttcctagatc	gtaaagatag	aggaaatcgt	ccattgtaat	3060
ctccggggca	aaggagatct	cttttggggc	tggatcactg	ctgggccttt	tggttcctag	3120
cgtgagccag	tgggcttttt	gctttggttg	gcttggttag	gccttagcaa	agctcttggg	3180
cttgagttga	gcttctcctt	tggggatgaa	gttcaacctg	tctgtttgct	gacttgttgt	3240
gtaecgctca	gctgtgctc	ttgcctctgt	aatagtggca	aatttcttgt	gtgcaactcc	3300
gggaacccg	ttgttgccg	cctttgtaca	acccagtc	tcgtatatac	cggcatgtgg	3360
accgttatac	acaacgtagt	agttgatatg	aggggtgtga	ataccgatt	ctgctctgag	3420
aggagcaact	gtgctgttaa	gctcagat	ttgtgggatt	ggaattaat	cgtcgagcgg	3480
ccgctcgacg	agcgcgccga	tatcgcgatc	gcccggggcg	gccatttaa	tgaattcgag	3540
ctcggatccc	aaacgcggcc	gcaagctata	acttctgata	gcatacatta	tacgaagtta	3600
ttcgactcta	gaggatccca	attcccatgc	atggagtcaa	agattcaaat	agaggacact	3660
tctcgaactc	ggcgcgtcga	ctcggccgtc	gagtacatgg	tcgataagaa	aaggcaattt	3720
gtagatgtta	attcccatct	tgaagaaat	atagttaa	tatttattga	taaaataaca	3780
agtcagggtat	tatagtccaa	gcaaaaacat	aaatttattg	atgcaagttt	aaattcagaa	3840
atatttcaat	aactgattat	atcagctggt	acattgccc	agatgaaaga	ctgagtgcga	3900
tattatgtgt	aatacataaa	ttgatgat	agctagctta	gctcatcggg	ggatcctaga	3960
cgcgtgagat	cagatctcgg	tgcggggcag	gaccggacgg	ggcggtagcg	gcaggtgaa	4020
gtccagctgc	cagaaaccca	cgtcatgcc	gttcccgtgc	ttgaagccgg	cgcggcgag	4080
catgccgcgg	ggggcatatc	cgagcgcctc	gtgcatgcgc	acgctcgggt	cgttgggcag	4140
ccgatgaca	gcgaccacgc	tcttgaagcc	ctgtgcctcc	agggacttca	gcaggtgggt	4200
gtagagcgtg	gagcccagtc	ccgtccgctg	gtggcggggg	gagacgtaca	cggctcgactc	4260
ggcgcgtccg	tcgtaggcgt	tgctgtcctt	ccagggggcc	gcgtaggcga	tgccggcgac	4320
ctcgcgcgtc	acctcgccga	cgagccaggg	atagcgtcc	cgcagacgga	cgaggtcgtc	4380
cgtccactcc	tgcggttctt	gcggctcggt	acggaagtgt	accgtgcttg	tctcgatgta	4440
gtggttgacg	atggtgcaga	cgcgcggcat	gtccgcctcg	gtggcacggc	ggatgtcggc	4500
cgggcgtcgt	tctgggtcca	ttgttcttct	ttactctttg	tgtgactgag	gtttggtcta	4560
gtgctttggt	catctatata	taagtataac	aacaatgaga	acaagctttg	gagtgatcgg	4620
aggggtctagg	atacatgaga	ttcaagtgga	ctaggatcta	caccgttggg	ttttgagtgt	4680
ggatatgtgt	gaggttaaatt	ttacttggta	acggccacaa	aggcctaagg	agaggtgttg	4740
agacccttat	ggccttgaac	cgtctggaata	atgccacgtg	gaagataatt	ccatgaatct	4800
tatcgttatc	tatgagtga	attgtgtgat	ggtggagtgg	tgcttgctca	ttttacttgc	4860
ctggtggact	tggccctttc	cttatgggga	atttatattt	tacttactat	agagctttca	4920
tacctttttt	ttaccttggg	tttagttaat	atataatggt	atgattcatg	aataaaaatg	4980
ggaaattttt	gaatttgcac	tgctaaatgc	ataagattag	gtgaaactgt	ggaatatata	5040
tttttttcat	ttaaaagcaa	aatttgcctt	ttactagaat	tataaatata	gaaaaatata	5100

taacattcaa	ataaaaaatga	aaataagaac	tttcaaaaaa	cagaactatg	tttaatgtgt	5160
aaagattagt	cgcacatcaa	gtcatctgtt	acaatatgtt	acaacaagtc	ataagcccaa	5220
caaagtttagc	acgtctaaat	aaactaaaga	gtccacgaaa	atattacaaa	tcataagccc	5280
aacaaagtta	ttgatcaaaa	aaaaaaaaacg	cccaacaaag	ctaaacaaag	tccaaaaaaa	5340
acttctcaag	tctccatctt	cctttatgaa	cattgaaaac	tatacacaaa	acaagtcaga	5400
taaatctctt	tctgggcctg	tcttcccaac	ctcctacatc	acttccctat	cggattgaat	5460
gttttacttg	taccttttcc	gttgcaatga	tattgatagt	atgtttgtga	aaactaatag	5520
ggttaacaat	cgaagtcacg	gaatatggat	ttgggtccaag	attttccgag	agctttctag	5580
tagaaagccc	atcaccagaa	atttactagt	aaaataaatc	accaattagg	tttcttatta	5640
tgtgccaaat	tcaatataat	tatagaggat	atttcaaatg	aaaacgtatg	aatgttatta	5700
gtaaatggtc	aggtaagaca	ttaaaaaaat	cctacgtcag	atattcaact	ttaaaaattc	5760
gatcagtggtg	gaattgtaca	aaaatttggg	atctactata	tatatataat	gctttacaac	5820
acttggattt	ttttttggag	gctggaattt	ttaatctaca	tatttgtttt	ggccatgcac	5880
caactcattg	tttagtga	tactttgatt	ttgtcaata	tatgtgttcg	tgtatatattg	5940
tataagaatt	tctttgacca	tatacacaca	cacatatata	tatatatata	tatattatat	6000
atcatgcaact	tttaattgaa	aaaataatat	atatatatat	agtgcatttt	ttctaacaac	6060
catatatgtt	gcgattgatc	tgcaaaaaata	ctgctagagt	aatgaaaaat	ataatctatt	6120
gctgaaatta	tctcagatgt	taagattttc	ttaaagttaa	ttctttcaaa	ttttagctaa	6180
aagtcttgta	ataactaaag	aataatacac	aatctcgacc	acggaaaaaa	aacacataat	6240
aaatttgaat	ttecgaccgcg	gtacccggaa	ttgggttata	attacctcag	gtcgaggaat	6300
taattcggta	cgtacctaat	aaacttcgtat	agcatacatt	atacgaagtt	atatggatct	6360
cgaggcatta	cggcattacg	gcactcgcca	gggtcccaat	tcgagcatgg	agccatttac	6420
aattgaatat	atcctgcgcg	cgctgccgct	ttgcacccgg	tggagcttgc	atgttgggtt	6480
ctacgcagaa	ctgagccggt	taggcagata	atttccattg	agaactgagc	catgtgcacc	6540
ttccccccaa	cacggtgagc	gacggggcaa	cggagtgatc	cacatgggac	ttttaaacat	6600
catccgtcgg	atggcggttcg	gagagaagca	gtcgatccgt	gagatcagcc	gacgcaccgg	6660
gcaggcgccg	aacacgatcg	caaagtattt	gaacgcaggt	acaatcgagc	cgacgttcac	6720
ggtagccgaa	cgaccaagca	agctagctta	gtaaagccct	cgctagattt	taatgccgat	6780
gttgcgatta	cttcgccaac	tattgcgata	acaagaaaaa	gccagccttt	catgatatat	6840
ctcccaattt	gtgtagggct	tattatgcac	gcttaaaaaat	aataaaagca	gacttgacct	6900
gatagtttgg	ctgtgagcaa	ttatgtgctt	agtgcattct	acgcttgagt	taagccgcgc	6960
cgcgaaagcg	cgctcgcttg	aacgaattgt	tagacattat	ttgcccacta	ccttgggtgat	7020
ctcgcccttc	acgtagtggg	caaattcttc	caactgatct	gcgcgcgagg	ccaagcgatc	7080
ttcttcttgt	ccaagataag	cctgtctagc	ttcaagtatg	acgggctgat	actgggcccgg	7140
caggcgctcc	attgcccagt	cggcagcgac	atccttcggc	gcgattttgc	cggttactgc	7200
gctgtaccaa	atgcccggaca	acgtaagcac	tacatttcgc	tcacgcgccag	cccagtcggg	7260
cggcgagttc	catagcgta	aggtttcatt	tagcgcctca	aatagatcct	gttcagggaac	7320
cggatcaaaag	agttcctccg	ccgctggacc	taccaaggca	acgctatggt	ctcttgcttt	7380
tgtcagcaag	atagccagat	caatgtcgat	cgtggctggc	tcgaagatac	ctgcaagaat	7440
gtcattgcgc	tgccattctc	caaattgcag	ttcgcgctta	gctggataac	gccacggaat	7500
gatgtcgtcg	tgcaacaaca	tggtgacttc	tacagcgcgg	agaatctcgc	tctctccagg	7560
ggaagccgaa	gtttccaaaa	ggtcggtgat	caaagctcgc	cgcgttggtt	catcaagcct	7620
tacggtcacc	gtaaccagca	aatcaatatc	actgtgtggc	ttcaggccgc	catccactgc	7680
ggagccgtac	aaatgtacgg	ccagcaacgt	cggtttcgaga	tggcgctcga	tgacgcacaac	7740
tacctctgat	agttgagtcg	atacttcggc	gatcacccgt	tccctcatga	tgtttaactt	7800
tgttttaggg	cgactgccct	gctgcgtaac	atcgttgctg	ctccataaca	tcaaacatcg	7860
accacggcg	taacgcgctt	gctgcttgga	tgcccagggc	atagactgta	ccccaaaaaa	7920
acagtcataa	caagccatga	aaaccgccac	tgccgcgtta	ccaccgctgc	gttcgggtcaa	7980
ggttctggac	cagttgcgtg	agcgcatacg	ctacttgcac	tacagcttac	gaaccgaaca	8040
ggcttatgtc	cactgggttc	gtgccttcac	ccgtttccac	gggtgtgcgtc	acccggcaac	8100
cttgggcagc	agcgaaagtcg	aggcatttct	gtcctggctg	gcgaacgagc	gcaagggttc	8160
ggtctccacg	catcgtcagg	cattggcggc	cttgctgttc	ttctacggca	agtgtgtgtc	8220
acggatctgc	cctggcttca	ggagatcgga	agacctcggc	cgtccggggc	cttgccgggtg	8280
gtgctgaccc	cggatgaagt	ctctagagct	ctagaggggt	cgcatcctcg	gttttctgga	8340
aggcgagcat	cgtttgttcg	cccagcttct	gtatggaacg	ggcatgcgga	tcagtgaggg	8400
tttgcaactg	cgggtcaagg	acttggattt	cgatcacggc	acgatcatcg	tcggggaggg	8460
caagggtctcc	aaggatcggg	ccttgatgtt	acccgagagc	ttggcaccca	gcctgcgcga	8520
gcagggatcg	atccaaaccc	tccgctgcta	tagtgcagtc	ggcttctgac	gttcagtgc	8580
gccgtcttct	gaaaacgaca	tgtcgcacaa	gtcctaagtt	acgcgacagg	ctgccgcctt	8640
gcccttttcc	tggcgttttc	ttgtcgcgtg	tttttagtcg	ataaagtaga	atacttgcca	8700
ctagaaccgg	agacattacg	ccatgaacaa	gagcgccgcc	gctggcctgc	tgggctatgc	8760

ccgcgtcagc	accgacgacc	aggacttgac	caaccaacgg	gccgaactgc	acgcggccgg	8820
ctgcaccaag	ctgttttccg	agaagatcac	cggcaccagg	cgcgaccgcc	cggagctggc	8880
caggatgctt	gaccacctac	gccctggcga	cgttgtgaca	gtgaccaggc	tagaccgcct	8940
ggcccgagc	accccgagcc	tactggacat	tgccgagcgc	atccaggagg	ccggcgccgg	9000
cctgcgtagc	ctggcagagc	cgtgggcccga	caccaccacg	ccggccggcc	gcattggtgtt	9060
gaccgtgttc	gccggcattg	ccgagttcga	gcgttcccta	atcatcgacc	gcaccgagag	9120
cgggcgcgag	gccgccaagg	cccaggcggt	gaagtttggc	ccccgccta	ccctcaccce	9180
ggcacagatc	gcgcacgccc	gcgagctgat	cgaccaggaa	ggccgcaccg	tgaaagaggc	9240
ggctgactg	cttggcgctg	atcgctcgac	cctgtaccgc	gcacttgagc	gcagcgagga	9300
agtgacgccc	accgaggcca	ggcgccgcgg	tgcttccgt	gaggacgcat	tgaccgagge	9360
cgacgcccctg	gcggccgcgg	agaatgaacg	ccaagaggaa	caagcatgaa	accgcaccag	9420
gacggccagg	acgaaccggt	tttcattacc	gaagagatcg	aggcgagat	gatcgccggc	9480
gggtacgtgt	tcgagccgcc	cgcgacgtc	tcaaccgtgc	ggctgcatga	aatcctggcc	9540
ggtttgtctg	atgccaaagt	ggcgccctgg	ccggccagct	tgcccgctga	agaaaccgag	9600
cgccgcgctc	taaaaagggtg	atgtgtattt	gagtaaaaca	gcttgcgctc	tgccgtcgct	9660
gcgtatatga	tgcatgagtg	aaataaaca	atacgcaagg	ggaacgcatg	aaggttatcg	9720
ctgtacttaa	ccgaaaggc	gggtcagga	agacgaccat	cgcaaccat	ctagcccgcg	9780
ccctgcaact	cgccggggcc	gatgttctgt	tagtcgattc	cgatccccag	ggcagtgccc	9840
gcgattgggc	ggccgtgccc	gaagatcaac	cgctaaccgt	tgccgcatc	gaccgcccga	9900
cgattgaccg	cgacgtgaag	gccatcgccc	ggcgcgactt	cgtagtgatc	gacggagcgc	9960
cccaggcgcc	ggacttggct	gtgtccgcga	tcaaggcagc	cgacttcgtg	ctgattccgg	10020
tgacgccaag	cccttacgac	atatgggcca	ccgcgcacct	ggtggagctg	gttaagcagc	10080
gcattgaggt	cacggatgga	aggctacaag	cgcccttctg	cggtgcgccc	gcgatcaaa	10140
gcacgcgcac	cgccgggtgag	ggtgcccagg	cgctggccgg	gtacgagctg	ccattctctg	10200
agtcccgtat	cacgcagcgc	gtgagctacc	caggcactgc	cgccgcggcc	acaaccgttc	10260
ttgaatcaga	acccgagggg	gacgtgccc	gcgaggtcca	ggcgctggcc	gctgaaatta	10320
aatcaaaact	catttgagtt	aatgaggtaa	agagaaaatg	agcaaaagca	caaacacgct	10380
aagtgcgggc	cgtccgagcg	cacgcagcag	caaggctgca	acgttggcca	gcctggcaga	10440
cacgccagcc	atgaagcggg	tcaactttca	ggttccggcg	gaggatcaca	ccaagctgaa	10500
gatgtacgcg	gtacgccaag	gcaagaccat	taccgagctg	ctatctgaat	acatcgcgca	10560
gctaccagag	taaatgagca	aatgaataaa	tgagttagtg	aatttttagc	gctaaaggag	10620
gcggcatgga	aaatcaagaa	caaccaggca	ccgacgccgt	ggaatgcccc	atgtgtggag	10680
gaacggggcg	ttggccaggc	gtaagcggct	gggttgtctg	ccggccctgc	aatggcactg	10740
gaacccccaa	gcccagggaa	tcggcggtgac	ggtcgcaaac	catccggccc	ggtacaaatc	10800
ggcgcgccgc	tggtgatga	cctggtggag	aaagtgaagg	ccgcgcaggc	cgcccagcgg	10860
caacgcatcg	aggcagaagc	acgccccggt	aatcgctggc	aagcgccgcg	tgatcgaatc	10920
cgaaagaat	cccggcaacc	gccggcagcc	ggtgcgccgt	cgattaggaa	gccgcccag	10980
ggcgacgagc	aaccagattt	tttcgttccg	atgctctatg	acgtgggcac	ccgcgatagt	11040
cgcgacatca	tggacgtggc	cgttttccgt	ctgtcgaagc	gtgaccgacg	agctggcgag	11100
gtgatccgct	acgagcttcc	agacgggcac	gtagaggttt	ccgcaggggc	ggccggcatg	11160
gccagtgtgt	gggattacga	cctggtactg	atggcggttt	cccatctaac	cgaatccatg	11220
aaacgatacc	gggaaggga	gggagacaag	cccgcccgcg	tggtccgctc	acacgttgcg	11280
gacgtactca	agttctgccc	gcgagccgat	ggcggaagc	agaaagacga	cctggtagaa	11340
acctgcattc	ggttaaaccac	cacgcacggt	gccatgcagc	gtacgaagaa	ggccaagaac	11400
ggccgcctgg	tgacgggtatc	cgagggtgaa	gccttgatta	gccgctacaa	gatcgtaaa	11460
agcgaaaccg	ggcgcccgga	gtacatcgag	atcgagctag	ctgattggat	gtaccgcgag	11520
atcacagaag	gcaagaacc	ggacgtgctg	acggttcacc	ccgattactt	tttgatcgat	11580
cccggcatcg	cggtttttct	ctaccgcctg	gcacgcgcg	ccgcaggcaa	ggcagaagcc	11640
agatgggtgt	tcaagacgat	ctacgaacgc	agtggcagcg	ccggagagtt	caagaagttc	11700
tgtttcaccg	tgccgaagct	gatcggttca	aatgacctgc	cggagtacga	tttgaaggag	11760
gaggcggggc	aggctggccc	gatcctagtc	atgcgctacc	gcaacctgat	cgaggggcga	11820
gcatccgcgc	ggttctaattg	tacggagcag	atgctagggc	aaattgccct	agcaggggaa	11880
aaaggctcga	aaggtctctt	tcctgtggat	agcagctaca	ttgggaacct	aaagccgtac	11940
attgggaacc	ggaacccgta	cattgggaac	ccaaagccgt	acattgggaa	ccggtcacac	12000
atgtaagtga	ctgatataaa	agagaaaaaa	ggcgattttt	ccgcctaaaa	ctctttaaaa	12060
cttattaaaa	ctcttaaaac	ccgcctggcc	tggtgcataac	tgcttgccca	gcgcacagcc	12120
gaagagctgc	aaaaagcgcc	tacccttcgg	tcgctgcgct	ccctacgccc	cgccgcttcg	12180
cgctggccta	tcgcggccgc	tgcccgctca	aaaatggctg	gcctacggcc	aggcaatcta	12240
ccaggcgccg	gacaagccgc	gccgtcgcca	ctcgaccgcc	ggcgcccaca	tcaaggcacc	12300
ctgcctcgcg	cgtttcgggtg	atgacgggtg	aaacctctga	cacatgcagc	tcccgagagc	12360
ggtcacagct	tgtctgtaag	cggatgccgg	gagcagacaa	gcccgctcagg	gcgcgtcagc	12420


```

gggtgttggc ggggtgtcggg ggcagccat gacccagtca cgtagcgata gcggagtgt 12480
tactggctta actatgcggc atcagagcag attgtactga gtagtcacca tatgcgggtgt 12540
gaaataccgc acagatg~cgt aaggagaaaa taccgcatca ggcgtcttc cgcttcctcg 12600
ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag cggatcagc tcaactcaaag 12660
gcggtaatac ggttatccac agaatcaggg gataacgcag gaaagaacat gtgagcaaaa 12720
ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc tggcgttttt ccataggctc 12780
cgccccctg acgagcatca caaaaatcga cgctcaagtc agaggtggcg aaacccgaca 12840
ggactataaa gataccaggc gtttccccct ggaagctccc tcgtgcgctc tcctgttccg 12900
accctgcgcg ttaccggata cctgtccgcc tttctccctt cgggaagcgt ggcgctttct 12960
catagctcac gctgtaggta tctcagttcg gtgtaggtcg ttcgctccaa gctgggctgt 13020
gtgcacgaac ccccggttca gcccgaccgc tgcgccttat ccgtaacta tcgtcttgag 13080
tccaacccgg taagacacga cttatcgcca ctggcagcag ccactggtaa caggattagc 13140
agagcgaggt atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac 13200
actagaagga cagtatttgg tatctgcgct ctgctgaagc cagttacctt cggaaaaaga 13260
gttggttagct cttgatccgg caaacaacc accgctggta gcggtggttt ttttgtttgc 13320
aagcagcaga ttacgcgcag aaaaaaagga tctcaagaag atccggaaaa cgcaagcgca 13380
aagagaaagc aggtagcttg cagtgggctt acatggcgat agctagactg ggcggtttta 13440
tggacagcaa gcgaaccgga attgcc 13466

```

<210> 10

<211> 31

<212> PRT

<213> Artificial Sequence

<220>

<223> Consensus sequence 1 of PARG protein

<220>

<221> MISC_FEATURE

<222> (2)..(2)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (8)..(10)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (14)..(17)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (19)..(19)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (27)..(29)

<223> X represents any amino acid

<400> 10

```

Leu Xaa Val Asp Phe Ala Asn Xaa Xaa Xaa Gly Gly Gly Xaa Xaa Xaa
1           5           10           15

```

```

Xaa Gly Xaa Val Gln Glu Glu Ile Arg Phe Xaa Xaa Xaa Pro Glu
20           25           30

```

<210> 11

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Consensus sequence 2 for PARG protein

<220>

<221> MISC_FEATURE

<222> (3)..(3)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (10)..(10)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (13)..(14)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (17)..(19)

<223> X represents any amino acid

<400> 11

Thr Gly Xaa Trp Gly Cys Gly Ala Phe Xaa Gly Asp Xaa Xaa Leu Lys
1 5 10 15

Xaa Xaa Xaa Gln
20

<210> 12

<211> 13

<212> PRT

<213> Artificial Sequence

<220>

<223> Consensus sequence 3 for PARG protein

<220>

<221> MISC_FEATURE

<222> (2)..(4)

<223> X represents any amino acid

<220>

<221> MISC_FEATURE

<222> (6)..(9)

<223> X represents any amino acid

<400> 12

Asp Xaa Xaa Xaa Arg Xaa Xaa Xaa Xaa Ala Ile Asp Ala
1 5 10

<210> 13

<211> 10

<212> PRT

<213> Artificial Sequence

<220>

<223> Consensus sequence 4 for PARG protein

<220>

<221> MISC_FEATURE

<222> (3)..(4)
 <223> X represents any amino acid

<220>
 <221> MISC_FEATURE
 <222> (7)..(8)
 <223> X represents any amino acid

<400> 13
 Arg Glu Xaa Xaa Lys Ala Xaa Xaa Gly Phe
 1 5 10

<210> 14
 <211> 11
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> conserved PARG region
 <220>
 <221> MISC_FEATURE
 <222> (2)..(5)
 <223> X represents any amino acid

<220>
 <221> MISC_FEATURE
 <222> (7)..(7)
 <223> X represents any amino acid

<400> 14
 Gly Xaa Xaa Xaa Ser Xaa Tyr Thr Gly Tyr
 1 5 10

<210> 15
 <211> 1530
 <212> DNA
 <213> Oryza sativa

<220>
 <221> CDS
 <222> (1)..(1530)
 <223>

<400> 15
 atg gag gcg cgc ggc gac ctg cgc tcg atc ctg ccc tac ctc ccc gtc 48
 Met Glu Ala Arg Gly Asp Leu Arg Ser Ile Leu Pro Tyr Leu Pro Val
 1 5 10 15
 gtg ctc cgc ggc ggc gcg ctc ttc tgg ccg ccg gcg gcg cag gag gcg 96
 Val Leu Arg Gly Gly Ala Leu Phe Trp Pro Pro Ala Ala Gln Glu Ala
 20 25 30
 ctc aag gcg ctg gcg ctg ggc ccc gac gtg agc cgc gtc tcc tcc ggc 144
 Leu Lys Ala Leu Ala Leu Gly Pro Asp Val Ser Arg Val Ser Ser Gly
 35 40 45
 gac gtc ctc gcc gac gcc ctc acc gac ctc cgc ctc gcg ctc aac ctc 192
 Asp Val Leu Ala Asp Ala Leu Thr Asp Leu Arg Leu Ala Leu Asn Leu
 50 55 60
 gac cca ctc ccg cgc cgc gcc gcc gag ggc ttc gcg ctc ttc ttc gac 240
 Asp Pro Leu Pro Arg Arg Ala Ala Glu Gly Phe Ala Leu Phe Phe Asp
 65 70 75 80
 gac ctc ctg tcg cgg gcg cag gcg cgg gac tgg ttc gac cac gtc gcc 288
 Asp Leu Leu Ser Arg Ala Gln Ala Arg Asp Trp Phe Asp His Val Ala

ttt ggt gga aac cct gaa atc aag agc atg att caa tgg att gct gca 1296
 Phe Gly Gly Asn Pro Glu Ile Lys Ser Met Ile Gln Trp Ile Ala Ala
 420 425 430
 tca cag gca ctc cga tct ttt att aac tac tac act ttt gag tcc gaa 1344
 Ser Gln Ala Leu Arg Ser Phe Ile Asn Tyr Tyr Thr Phe Glu Ser Glu
 435 440 445
 tca ctg aaa aga tta gaa gag gtg acc cag tgg ata ttg cgc cat agg 1392
 Ser Leu Lys Arg Leu Glu Glu Val Thr Gln Trp Ile Leu Arg His Arg
 450 455 460
 tgg acg gtt ggc gag ttg tgg gac atg ctt gtg gag tat tca tcc cag 1440
 Trp Thr Val Gly Glu Leu Trp Asp Met Leu Val Glu Tyr Ser Ser Gln
 465 470 475 480
 agg cta aga gga gac acc aat gag ggc ttt tta aca tgg cta ctt ccc 1488
 Arg Leu Arg Gly Asp Thr Asn Glu Gly Phe Leu Thr Trp Leu Leu Pro
 485 490 495
 aag gac atc ccc aat ggt gat gta gat tac atg tgt gaa tag 1530
 Lys Asp Ile Pro Asn Gly Asp Val Asp Tyr Met Cys Glu
 500 505

<210> 16

<211> 509

<212> PRT

<213> Oryza sativa

<400> 16

Met Glu Ala Arg Gly Asp Leu Arg Ser Ile Leu Pro Tyr Leu Pro Val
 1 5 10 15

Val Leu Arg Gly Gly Ala Leu Phe Trp Pro Pro Ala Ala Gln Glu Ala
 20 25 30

Leu Lys Ala Leu Ala Leu Gly Pro Asp Val Ser Arg Val Ser Ser Gly
 35 40 45

Asp Val Leu Ala Asp Ala Leu Thr Asp Leu Arg Leu Ala Leu Asn Leu
 50 55 60

Asp Pro Leu Pro Arg Arg Ala Ala Glu Gly Phe Ala Leu Phe Phe Asp
 65 70 75 80

Asp Leu Leu Ser Arg Ala Gln Ala Arg Asp Trp Phe Asp His Val Ala
 85 90 95

Pro Ser Leu Ala Arg Leu Leu Leu Arg Leu Pro Thr Leu Leu Glu Gly
 100 105 110

His Tyr Arg Ala Ala Gly Asp Glu Ala Arg Gly Leu Arg Ile Leu Ser
 115 120 125

Ser Gln Asp Ala Gly Leu Val Leu Leu Ser Gln Glu Leu Ala Ala Ala
 130 135 140

Leu Leu Ala Cys Ala Leu Phe Cys Leu Phe Pro Thr Ala Asp Arg Ala
 145 150 155 160

Glu Ala Cys Leu Pro Ala Ile Asn Phe Asp Ser Leu Phe Ala Ala Leu
 165 170 175

Cys Tyr Asn Ser Arg Gln Ser Gln Glu Gln Lys Val Arg Cys Leu Val
 180 185 190

His Tyr Phe Asp Arg Val Thr Ala Ser Thr Pro Thr Gly Ser Val Ser
 195 200 205
 Phe Glu Arg Lys Val Leu Pro Arg Arg Pro Glu Ser Asp Gly Ile Thr
 210 215 220
 Tyr Pro Asp Met Asp Thr Trp Met Lys Ser Gly Val Pro Leu Cys Thr
 225 230 235 240
 Phe Arg Val Phe Ser Ser Gly Leu Ile Glu Asp Glu Glu Gln Glu Ala
 245 250 255
 Leu Glu Val Asp Phe Ala Asn Arg Tyr Leu Gly Gly Gly Ala Leu Ser
 260 265 270
 Arg Gly Cys Val Gln Glu Glu Ile Arg Phe Met Ile Asn Pro Glu Leu
 275 280 285
 Ile Val Gly Met Leu Phe Met Val Ser Met Glu Asp Asn Glu Ala Ile
 290 295 300
 Glu Ile Val Gly Ala Glu Arg Phe Ser Gln Tyr Met Gly Tyr Gly Ser
 305 310 315 320
 Ser Phe Arg Phe Thr Gly Asp Tyr Leu Asp Ser Lys Pro Phe Asp Ala
 325 330 335
 Met Gly Arg Arg Lys Thr Arg Ile Val Ala Ile Asp Ala Leu Asp Cys
 340 345 350
 Pro Thr Arg Leu Gln Phe Glu Ser Ser Gly Leu Leu Arg Glu Val Asn
 355 360 365
 Lys Ala Phe Cys Gly Phe Leu Asp Gln Ser Asn His Gln Leu Cys Ala
 370 375 380
 Lys Leu Val Gln Asp Leu Asn Thr Lys Asp Asn Cys Pro Ser Val Ile
 385 390 395 400
 Pro Asp Glu Cys Ile Gly Val Ser Thr Gly Asn Trp Gly Cys Gly Ala
 405 410 415
 Phe Gly Gly Asn Pro Glu Ile Lys Ser Met Ile Gln Trp Ile Ala Ala
 420 425 430
 Ser Gln Ala Leu Arg Ser Phe Ile Asn Tyr Tyr Thr Phe Glu Ser Glu
 435 440 445
 Ser Leu Lys Arg Leu Glu Glu Val Thr Gln Trp Ile Leu Arg His Arg
 450 455 460
 Trp Thr Val Gly Glu Leu Trp Asp Met Leu Val Glu Tyr Ser Ser Gln
 465 470 475 480
 Arg Leu Arg Gly Asp Thr Asn Glu Gly Phe Leu Thr Trp Leu Leu Pro
 485 490 495
 Lys Asp Ile Pro Asn Gly Asp Val Asp Tyr Met Cys Glu
 500 505

<210> 17

<211> 25
<212> DNA
<213> Artificial Sequence

<220>
<223> degenerate oligonucleotide primer PG1
<400> 17
atgtbccaca rmtckccrac mgtcc

25

<210> 18
<211> 28
<212> DNA
<213> Artificial Sequence

<220>
<223> degenerate oligonucleotide primer PG2
<400> 18
gggtytccwc caaaarcmmc rcawcccc

28

<210> 19
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> degenerate oligonucleotide primer PG3
<400> 19
gctatagaaa twgtyggtgy rgaaag

26

<210> 20
<211> 26
<212> DNA
<213> Artificial Sequence

<220>
<223> degenerate oligonucleotide primer PG4
<400> 20
agrggstgyg trcaggarga ratmcg

26

<210> 21
<211> 23
<212> DNA
<213> Artificial Sequence

<220>
<223> degenerate oligonucleotide primer PG5
<220>
<221> misc_feature
<222> (18)..(18)
<223> n=any nucleotide

<400> 21
atggargaya aygargcnat hga

23

<210> 22
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> degenerate oligonucleotide primer PG6

<400> 22
ccaytgdagc atrctytttda gytc

24

<210> 23
<211> 603
<212> DNA
<213> zea mays

<400> 23
tagggctgtg tgcaggagga aatccgcttc atgataaacc ccgaattgat tgtgggtatg 60
ctattcttgt cttgtatgga agataacgag gctatagaaa tctttggtgc agaacgggtc 120
tcacagtata tgggttatgg ttcctccttt cgctttgttg gtgactattt agataccaaa 180
ccctttgatt cgatgggcag acggagaact aggattgttg ctatcgatgc tttggactgt 240
ccagctagggt tacactatga atctggctgt ctccctaaggg aagtgaacaa ggcattttgt 300
ggatttttcg atcaatcgaa acaccatctc tatgcgaagc ttttcagga tttgcacaac 360
aaggatgact tttcaagcat caattccagt gagtacgtag gagtttcaac aggaaactgg 420
ggttgtgggtg cttttgtggg aaaccctgaa atcaagagca tgattcagtg gattgctgca 480
tcacaggctc ttcgcccttt tgtaattac tacacttttg agaacgtgtc tctgcaaaga 540
ttagaggagg tgatccagtg gatacggctt catggctgga ctgtcggcga gctgtggaac 600
ata 603

<210> 24
<211> 12987
<212> DNA
<213> Artificial sequence

<220>
<223> T-DNA vector comprising a chimeric ParG expression reducing gene
<220>
<221> misc_feature
<222> (1)..(25)
<223> complement of Left T-DNA border

<220>
<221> misc_feature
<222> (58)..(318)
<223> complement of 3' nos

<220>
<221> misc_feature
<222> (337)..(888)
<223> bar coding region

<220>
<221> misc_feature
<222> (889)..(1721)
<223> 35S promoter

<220>
<221> misc_feature
<222> (1728)..(3123)
<223> 35S promoter

<220>
<221> misc_feature
<222> (3133)..(3311)
<223> part of ParG homologue of Zea mays

<220>
<221> misc_feature
<222> (3344)..(4085)

<223> Pdk intron

<220>

<221> misc_feature

<222> (4119)..(4297)

<223> part of ParG homologue of Zea mays (inverted)

<220>

<221> misc_feature

<222> (4310)..(5020)

<223> 3' OCS

<220>

<221> misc_feature

<222> (5066)..(5042)

<223> Right T-DNA border

<400> 24

cggcaggata	tattcaattg	taaatggctc	catggcgatc	gctctagagg	atcttcccga	60
tctagtaaca	tagatgacac	cgcgcgcgat	aattttatcct	agtttgccgcg	ctatatatttg	120
ttttctatcg	cgtattaaat	gtataattgc	gggactctaa	tcataaaaaac	ccatctcata	180
aataacgtca	tgcattacat	gttaattatt	acatgcttaa	cgtaattcaa	cagaaattat	240
atgataatca	tgcgaagacc	ggcaacagga	ttcaatctta	agaaacttta	ttgccaaatg	300
tttgaacgat	ctgcttcgga	tcctagacgc	gtgagatcag	atctcgggtga	cgggcaggac	360
cggacggggc	ggtagccgga	ggctgaagtc	cagctgccag	aaaccacagt	catgccagtt	420
cccgtgcttg	aagccggccg	cccgcagcat	gccgcggggg	gcataaccga	gcgcctcgtg	480
catgcgcacg	ctcgggtcgt	tgggcagccc	gatgacagcg	accacgctct	tgaagccctg	540
tgctccagg	gacttcagca	ggtgggtgta	gagcgtggag	cccagtcctg	tccgctgggtg	600
gcggggggag	acgtacacgg	tcgactcggc	cgtccagtcg	taggcgttgc	gtgccttcca	660
ggggcccgcg	taggcgatgc	cggcgacctc	gccgtccacc	tcggcgacga	gccagggata	720
gcgctccgcg	agacggacga	ggctcgtccg	ccactcctgc	ggttcctgcg	gctcggtagc	780
gaagttgacc	gtgcttgtct	cgatgtagt	gttgacgatg	gtgcagaccg	ccggcatgtc	840
cgctcgggtg	gcacggcgga	tgtcggccgg	gcgtcgttct	gggtccatgg	ttatagagag	900
agagatagat	ttatagagag	agactggtga	tttcagcgtg	tcctctccaa	atgaaatgaa	960
cttccttata	tagaggaagg	gtccttcgaa	ggatagtggtg	attgtgcgtc	atcccttacg	1020
tcagtggaga	tgccacatca	atccacttgc	tttgaagacg	tgggtggaac	gtcttctttt	1080
tccacgatgc	gaatccttaga	ctcctgtggg	tttttgggac	cactgtcggc	agaggcatct	1140
tgaatgatag	cctttccttt	atcgcaatga	tggcatttgt	aggagccacc	ttccttttct	1200
actgtccttt	cgatgaagt	acagatagct	gggcaatgga	atccgaggag	gtttcccgaa	1260
attatccttt	gttgaaaagt	ctcaatagcc	ccttgggtctt	ctgagactgt	atctttgaca	1320
tttttggagt	agaccagagt	gtcgtgctcc	accatggtga	cgaagatttt	cttcttgtca	1380
ttgagtcgta	aaagactctg	tatgaactgt	tcgccagtct	tcacggcgag	ttctgttaga	1440
tcctcgattt	gaatccttaga	ctccatgcat	ggccttagat	tcagtaggaa	ctaccttttt	1500
agagactcca	atctctatta	cttgcccttg	tttatgaagc	aagccttgaa	tcgtccatac	1560
tggaatagta	cttctgatct	tgagaaatat	gtctttctct	gtgttcttga	tgcaattagt	1620
cctgaatctt	ttgactgcat	ctttaaccct	ccttgggaagg	tatttgatct	cctggagatt	1680
gttactcggg	tagatcgtct	tgatgagacc	tgctgcgtag	gaacgcggcc	gcgtatacgt	1740
atcagatatct	tcgaattcga	gctcgtcgag	cggccgctcg	acgaattaat	tccaatccca	1800
caaaaatctg	agcttaacag	cacagttgct	cctctcagag	cagaatcggg	tattcaacac	1860
cctcatatca	actactacgt	tgtgtataac	gggtccacatg	ccggtatata	cgatgactgg	1920
ggttgtacaa	aggcggcaac	aaacggcggt	cccggagttg	cacacaagaa	atttgccact	1980
attacagagg	caagagcagc	agctgacgcg	tacacaacaa	gtcagcaaac	agacaggttg	2040
aacttcaccc	ccaaaggaga	agctcaactc	aagcccaaga	gctttgctaa	ggccctaaca	2100
agccccacca	agcaaaaagc	ccactggctc	acgctaggaa	ccaaaaggcc	cagcagtgat	2160
ccagcccaaa	aagagatctc	tttggcccg	gagattacaa	tggacgattt	cctctatctt	2220
tacgatctag	gaagggaagt	cgaaggtgaa	ggtagcagca	ctatgttcac	cactgataat	2280
gagaagggtta	gcctcttcaa	tttcagaaag	aatgctgacc	cacagatggt	tagagaggcc	2340
tacgcagcag	gtctcatcaa	gacgatctac	ccgagtaaca	atctccagga	gatcaaatat	2400
cttcccaaga	agggttaaaga	tgcagtcaaa	agattcagga	ctaattgcat	caagaacaca	2460
gagaaagaca	tatttctcaa	gatcagaagt	actattccag	tatggacgat	tcaaggcttg	2520
cttcataaac	caaggcaagt	aatagagatt	ggagtctcta	aaaaggtagt	tcctactgaa	2580

tctaaggcca	tgcattggagt	ctaagattca	aatcgaggat	ctaacagaac	tcgccgtgaa	2640
gactggcgaa	cagttcatat	agagtctttt	acgactcaat	gacaagaaga	aaatcttcgt	2700
caacatgggtg	gagcacgaca	ctctgggtcta	ctccaaaaat	gtcaaagata	cagtctcaga	2760
agaccaaagg	gctattgaga	cttttcaaca	aaggataatt	tcgggaaacc	tcctcggatt	2820
ccattgcccc	gctatctgtc	acttcatcga	aaggacagta	gaaaaggaag	gtggctccta	2880
caaatgccat	cattgcgata	aaggaaaggc	tatcattcaa	gatctctctg	ccgacagtgg	2940
tcccaaagat	ggacccccac	ccacgaggag	catcgtggaa	aaagaagacg	ttccaaccac	3000
gtcttcaaag	caagtggatt	gatgtgacat	ctccactgac	gtaagggatg	acgcacaatc	3060
ccactatcct	tcgcaagacc	cttctcttat	ataaggaagt	tcatttcatt	tggagaggac	3120
acgctcgagc	ccgaattgat	tgtgggtatg	ctattcttgt	cttgtatgga	agataacgag	3180
gctatagaaa	tctttggtgc	agaacggttc	tcacagtata	tgggttatgg	ttcctccttt	3240
cgctttgttg	gtgactattt	agataccaaa	ccctttgatt	cgatgggcag	acggagaact	3300
aggattgtgg	cggtaaccca	gcttggtaag	gaaataatta	ttttcttttt	tccttttagt	3360
ataaaaatagt	taagtgatgt	taattagtat	gattataata	atatagttgt	tataattgtg	3420
aaaaaataat	ttataaatat	attgtttaca	taaacaacat	agtaatgtaa	aaaaatatga	3480
caagtgatgt	gtaagacgaa	gaagataaaa	gttgagagta	agtataattat	ttttaatgaa	3540
tttgatcgaa	catgtaagat	gatatactag	cattaatatt	tgttttaatc	ataatagtaa	3600
ttctagctgg	tttgatgaat	taaataatcaa	tgataaaaata	ctatagtaaa	aataagaata	3660
aataaattaa	aataatattt	ttttatgatt	aatagtttat	tatataatta	aatatctata	3720
ccattactaa	atatttttagt	ttaaaagtta	ataaatattt	tgttagaaat	tccaatctgc	3780
ttgtaattta	tcaataaaca	aaatattaaa	taacaagcta	aagtaacaaa	taatatcaaa	3840
ctaatagaaa	cagtaatcta	atgtaacaaa	acataatcta	atgctaatat	aacaaagcgc	3900
aagatctatc	attttatata	gtattatttt	caatcaacat	tcttattaat	ttctaaataa	3960
tactttagtg	tttattaact	tctaaatgga	ttgactatta	attaaatgaa	ttagtcgaa	4020
atgaataaac	aaggtaacat	gtatagtcac	gtcatttgtg	tatcattgat	cttacatttg	4080
gattgattac	agttgggaag	ctgggttcga	aatcgatagc	cacaatccta	gttctccgtc	4140
tgcccatcga	atcaaagggt	ttggatctca	aatagtcacc	aacaaagcga	aaggagggaac	4200
cataacccat	atactgtgag	aaccgttctg	caccaaagat	ttctatagcc	tcgttatctt	4260
ccatacaaga	caagaatagc	ataccacaaa	tcaattcggg	tctagagtcc	tgctttaatg	4320
agatatgcga	gacgcctatg	atcgcattgat	atttgccttc	aattctgttg	tgacgttgt	4380
aaaaaacctg	agcatgtgta	gtccagatcc	ttaccggcgg	tttcgggtta	ttctaatagaa	4440
tatatcaccc	gttactatcg	tatttttatg	aataaatattc	ttcgttcaat	ttactgattg	4500
taccctacta	cttatatgta	caatattaaa	atgaaaacaa	tatattgtgc	tgaatagggt	4560
tatagcgaca	tctatgatag	agcgccacaa	taacaaacaa	ttgcgtttta	ttattacaaa	4620
tccaatttta	aaaaaagcgg	cagaaccggt	caaaccctaaa	agactgatta	cataaatctt	4680
attcaaat	caaaaggccc	caggggctag	tactacgac	acaccgagcg	gcgaactaat	4740
aacgttcact	gaagggaact	cgggttcccc	ccgggcgcgc	atgggtgaga	ttccttgaag	4800
ttgagtattg	gcggtccgct	ctaccgaaag	ttaccgggcac	cattcaaccc	ggtccagcac	4860
ggcgcccggg	taaccgactt	gctgccccga	gaattatgca	gcattttttt	ggtgtatgtg	4920
ggccccaaat	gaagtgcagg	tcaaaccctg	acagtgcaga	caaactcgtt	ggcggggtcca	4980
gggcgaattt	tgcgacaata	tgtcgaggct	cagcaggacc	tgccaggtcga	cgcccgagta	5040
ctggcaggat	atataccgtt	gtaatttgtc	gcgtgtgaat	aagtcgctgt	gtatgtttgt	5100
ttgattgttt	ctgttggagt	tcacccgcat	agtcggctag	attgatttag	atttcgttgt	5160
ccctgatgaa	ctgcccaggg	gaagccatct	tgagcgcgga	atgggaatgg	atttcgttgt	5220
acaacgagac	gacagaacac	ccacgggacc	gagcttcgat	cgagcatcaa	atgaaactgc	5280
aattttattca	tatcaggatt	atcaatacca	tatttttgaa	aaagccgttt	ctgtaatgaa	5340
ggagaaaact	caccgaggca	gttccatagg	atggcaagat	cctgggtatcg	gtctgcgatt	5400
ccgactcgtc	caacatcaat	acaacctatt	aatttccct	cgtcaaaaat	aaggttatca	5460
agtgagaaat	caccatgagt	gcgactgaa	ttccggtgaga	atggcaaaag	tttatgcatt	5520
tctttccaga	ctgtttcaac	aggccagcca	ttacgctcgt	catcaaaatc	actcgcatca	5580
accaaaccgt	tattcattcg	tgattgcgcc	tgagcgagac	gaaatacgcc	gctgttaaaa	5640
ggacaattac	aaacaggaat	cgaatgcaac	cggcgagga	acactgccag	cgcatcaaca	5700
atattttcac	ctgaatcagg	atattcttct	aatacctgga	atgctgtttt	tcgggggatc	5760
gcagtgggtga	gtaacctgca	atcatcagga	gtacggataa	aatgcttgat	ggtcgggaaga	5820
ggcataaatt	ccgtcagcca	gtttagtctg	accatctcat	ctgtaacatc	attggcaacg	5880
ctacctttgc	catgttttcag	aaacaactct	ggcgcatcgg	gcttcccata	caatcgatag	5940
attgtcgcac	ctgattgccc	gacattatcc	gaatctggca	attccgggtc	gcttgctgtc	6000
cataaaaccg	ccagtcctag	ctatcgccat	gtaagccac	tgcaagctac	ctgctttctc	6060
tttgcgcttg	cgttttccgg	atcttcttga	gatccttttt	ttctgcgcgt	aatctgctgc	6120
ttgcaaacaa	aaaaaccacc	gctaccagcg	gtggttttgt	tgccggatca	agagctacca	6180
actctttttc	cgaaggtaac	tggcttcagc	agagcgcaga	taccaaatac	tgctcttcta	6240

gtgtagccgt	agttaggcca	ccacttcaag	aactctgtag	caccgcctac	atacctcgct	6300
ctgctaattc	tgttaccagt	ggctgctgcc	agtggcgata	agtcgtgtct	taccgggttg	6360
gactcaagac	gatagtacc	ggataaggcg	cagcggtcgg	gctgaacggg	gggttcgtgc	6420
acacagccca	gcttggagcg	aacgacctac	accgaactga	gatacctaca	gcgtgagcta	6480
tgagaaagcg	ccacgcttcc	cgaagggaga	aaggcggaca	ggatcccggt	aagcggcagg	6540
gtcggaaacag	gagagcgcac	gagggagctt	ccagggggaa	acgcctggta	tctttatagt	6600
cctgtcgggt	ttcgccacct	ctgacttgag	cgtcgatttt	tgtgatgtc	gtcagggggg	6660
cggagcctat	ggaaaaacgc	cagcaacggc	gcctttttac	ggttcctggc	cttttgcctg	6720
ccttttgctc	acatgttctt	tcctgcgtta	tcccctgatt	ctgtggataa	ccgtattacc	6780
gccttttgagt	gagctgatac	cgctcgccgc	agccgaacga	ccgagcgag	cgagtcagtg	6840
agcgaggaag	cggaagagcg	cctgatgcgg	tattttctcc	ttacgcattc	gtgcggtatt	6900
tcacaccgca	tatggtgcac	tctcagtaca	atctgctctg	atgccgcata	gttaagccag	6960
tatacactcc	gctatcgcta	ctgactggg	tcactggctgc	gccccgacac	ccgccaacac	7020
ccgctgacgc	gcccgtacgg	gcttgtctgc	tcccggcatc	cgcttacaga	caagctgtga	7080
ccgtctccgg	gagctgcatt	tgtagaggt	tttaccgctc	atcaccgaaa	cgcgcgaggc	7140
aggggtccct	gatgtgggcg	ccggcggtcg	agtggcgacg	cgcgcgcttg	tccgcgccct	7200
ggtagattgc	ctggccgtag	gccagccatt	tttgagcggc	cagcgccgcg	gataggccga	7260
cgcgaaagcg	cggggcgtag	ggagcgcagc	gaccgaagg	taggcgcttt	ttgcagctct	7320
tcggctgtgc	gctggccaga	cagttatgca	caggccaggc	gggttttaag	agttttaata	7380
agttttaaa	agtttttaggc	ggaaaaatcg	ccttttttct	cttttatatc	agtcacttac	7440
atgtgtgacc	ggttcccaat	gtacggcttt	gggttcccaa	tgtacgggtt	ccggttccca	7500
atgtacggct	ttgggttccc	aatgtacgtg	ctatccacag	gaaagagacc	ttttcgacct	7560
ttttccctcg	ctagggcaat	ttgccctagc	atctgctccg	tacattagga	accggcggat	7620
gcttcgcccc	cgatcagggt	gcggtagcgc	atgactagga	tcgggcccagc	ctgccccgcc	7680
tccttccttca	aatcgctactc	cggcaggtca	tttgaccgga	tcagcttgcg	cacggtgaaa	7740
cagaacttct	tgaactctcc	ggcgctgcca	ctgcgttcgt	agatcgtctt	gaacaaccat	7800
ctggcttctg	ccttgccctgc	ggcgcgcggt	gccagggcgt	agagaaaaacg	gccgatgccg	7860
ggatcgatca	aaaagtaatc	gggggtgaacc	gtcagcacgt	ccgggttctt	gccttctgtg	7920
atctcgcggg	acatccaatc	agctagctcg	atctcgatgt	actccggccg	cccggtttctg	7980
ctctttacga	tcttgtagcg	gctaatacaag	gcttcaccct	cggataaccgt	caccaggcgg	8040
ccgttcttgg	ccttcttctg	acgctgcattg	gcaacgtgcg	tgggtgttaa	ccgaatgcag	8100
gtttctacca	ggtcgtcttt	ctgctttccg	ccatcggtc	gccggcagaa	cttgagtacg	8160
tcgcgaacgt	gtggacggaa	cacgcggccg	ggcttgtctc	ccttcccttc	ccggtatcgg	8220
ttcatggatt	cggttagatg	ggaaaccgcc	atcagtacca	ggctcgtaatc	ccacacactg	8280
gccatgcggg	ccggccctgc	ggaaacctct	acgtgcccg	ctggaagctc	gtagcggatc	8340
acctcgccag	ctcgtcggtc	acgcttcgac	agacggaaaa	cggccacgtc	catgatgctg	8400
cgactatcgc	gggtgcccac	gtcatagac	atcggaacga	aaaaatctgg	ttgctcgtcg	8460
cccttgggcg	gcttccctaat	cgacggcgca	ccggctgccg	gcgggttgccg	ggattctttg	8520
cggtatcgat	cagcggccgc	ttgccacgat	tcaccggggc	gtgcttctgc	ctcgatgcgt	8580
tgccgctggg	cggcctgcgc	ggccttcaac	ttctccacca	ggctcatcacc	cagcgcccg	8640
ccgatttgta	ccgggcccga	tggtttgcga	ccgtcacgcc	gattctcctg	gcttgggggt	8700
tcacgtgcca	ttgcaggggc	ggcagacaac	ccagccgctt	acgcctggcc	aaccgcccgt	8760
tcctccacac	atggggcatt	ccacggcgctc	gggtgcctggt	tgttcttgat	tttccatgcc	8820
gcctccttta	gccgctaaaa	ttcatctact	catttattca	tttgcctatt	tactctggta	8880
gctgcgcgat	gtattcagat	agcagctcgg	taatggtctt	gccttggcgt	accgcgtaca	8940
tcttcagctt	ggtgtgatcc	tccgcggcca	actgaaagtt	gacccgcttc	atggctggcg	9000
tgtctgccag	gctggccaac	ggtgcagcct	tgctgctgcg	tgcgctcgga	cggccggcac	9060
ttagcgtggt	tggtcttttg	ctcattttct	ctttacctca	ttactcaaaa	tgagttttga	9120
tttaatttca	gcggccacgg	cctggacctc	gcgggcagcg	tcgccctcgg	gttctgattc	9180
aagaacgggt	gtgcggcgcg	cggcagtgcc	tgggtagctc	acgcgctgcg	tgatacggga	9240
ctcaagaatg	ggcagctcgt	acccggccag	cgctcggcca	acctoaccgc	cgatgcgcgt	9300
gcctttgatc	gcccgcgaca	cgacaaaggc	cgctttagtc	cttccatccg	tgacctcaat	9360
gcgctgctta	accagctcca	ccaggtcggc	gggtggccat	atgtcgtaag	ggcttggctg	9420
caccggaatc	agcacgaagt	cggctgcctt	gatcgcgga	acagccaagt	ccgcgcgctg	9480
ggcgctccg	tcgatcacta	cgaagtccgc	ccggccgatg	gccttcacgt	cgcgggtcaat	9540
cgctcggcg	tcgatgccga	caacgggttag	cgggtgatct	tcccgacagg	ccgcccacac	9600
gcgggactg	ccctggggat	cgggaatcgac	taacagaaca	tcggcccccg	cgagttgcag	9660
ggcgcgggct	agatgggttg	cgatggctcg	cttgccctgac	ccgcctttct	ggttaagtac	9720
agcgataacc	ttcatgcgtt	ccccctgcgt	atttgtttat	ttactcatcg	catcataac	9780
gcagcgaccg	catgacgcaa	gctgttttac	tcaaatacac	atcacctttt	tagacggcgg	9840
cgctcgggtt	cttcagcggc	caagctggcc	ggccaggccg	ccagcttgcc	atcagacaaa	9900

```

ccggccagga tttcatgcag ccgcacgggt gagacgtgcg cgggcggctc gaacacgtac 9960
ccggcccgga tcatctccgc ctgcattctt tcggtaaatga aaaacgggtc gtcctggccg 10020
tcctggtgcg gtttcaTgct tgttcctctt ggcggttcatt ctccggcgcc gccaggggcg 10080
cggcctcggt caatgcgtcc tcacggaagg caccgcgcgc cctggcctcg gtgggcgtca 10140
cttcctcgct gcgctcaagt gcgcggtaca gggtcgagcg atgcacgcca agcagtgcag 10200
ccgcctcttt cacggtgcgg ccttcctggt cgatcagctc gcggcggtgc gcgatctgtg 10260
ccgggggtgag ggtagggcgg gggccaaact tcacgcctcg ggccttgccg gcctcgccg 10320
cgctccgggt gcggtcgatg attagggaa cgtcgaactc ggcaatgccg gcgaacacgg 10380
tcaaacacat gcggccggcc ggcggtggtg tgcggccca cggctctgcc aggtacgca 10440
ggcccgccgc ggcctctggt atgcgctcgg caatgtccag taggtcgccg gtgctgccc 10500
ccaggcggtc tagcctgggt actgtcaca cgtcgccagg gcgtagggtg tcaagcatcc 10560
tgcccgagct cgggcgggtc gcctggtgc cggatctctt ctcgaaaac agcttggtgc 10620
agccggccgc gtgcagtctc gcccggtggt tggtaagtc ctggctcgct gtgctgacgc 10680
gggcatagcc cagcaggcca gcggcgccgc tcttggtcat ggcgtaagt ctccggttct 10740
agtcgcaagt attctacttt atgcgactaa aacacgcgac aagaaaacgc caggaagagg 10800
gcaggcgccg agcctgtcgc gtaacttagg acttggtcga catgtcgttt tcagaagacg 10860
gctgcaactga acgtcagaag ccgactgcac tatagcagcg gagggttggt atcgatccct 10920
gctcgccgag cgtgggtgcc aagctctcgg gtaacatcaa ggcccgatcc ttggagccct 10980
tgccctcccg cacgatgac gtgccgtgat cgaatccag atccttgacc cgcagttgca 11040
aaccctcact gatccgcagt cccgttccat acagaagctg ggcaacaaa cgatgctcgc 11100
cttcagaaaa accgaggatg cgaaccactt catccggggt cagcaccacc ggcaagccgc 11160
cggacggccg aggtcttcct atctcctgaa gccaggcgag atccgtgcac agcacttgcc 11220
gtagaagaac agcaaggccg ccaatgcctg acgatgcgtg gagaccgaaa cctgcccgtc 11280
gttcgcccag caggacagaa atgcctcgac ttgctgctg cccaaggttg ccgggtgacg 11340
cacaccgtgg aaacgggatg aggcacgaac ccagtggaac taagcctggt cggttcgtaa 11400
gctgtaatgc aagtagcgta tgcgctcacg caactggtcc agaacttga ccgaacgcag 11460
cgggtgtaac ggcgcagtgg cggtttctat ggcttggtat gactgttttt ttggggtaca 11520
gtctatgcct cgggcatcca agcagcaagc gcgttacgcc gtgggtcgat gtttgatgtt 11580
atggagcagc aacgatgtta cgcagcaggg cagtcgccct aaacaaaagt taaacatcat 11640
gaggaagcgc gtgatcccg aagtatcgac ttaactatca gaggtagttg gcgtcatcga 11700
gcgccatctc gaaccgacgt tgctggccgt acatttgtac ggctccgcag tggatggccg 11760
cctgaagcca cacagtgata ttgatttgcg ggttacgggt accgtaaggc ttgatgaaac 11820
aacgcggcga gctttgatca acgacctttt ggaacttcg gcttccccct gagagagcga 11880
gattctccgc gctgtagaag tcaccattgt tgtgcacgac gacatcattc cgtggcggtta 11940
tccagctaag cgcgaactgc aatttggaag atggcagcgc aatgacattc ttgcaggat 12000
cttcgagcca gcacgatcg acattgatct ggctatcttg ctgacaaaag caagagaaca 12060
tagcgttgcc ttggtaggtc cagcggcgga ggaactcttt gatccgggtc ctgaacagga 12120
tctatttgag gcgctaaatg aaaccttaac gctatggaac tcgccgcccg actgggctgg 12180
cgatgagcga aatgtagtgc ttacgttgct ccgcatttgg tacagcgag taaccggcaa 12240
aatcgcccg aaggatgtcg ctgccgactg ggcaatggag gcgctgccgg cccagtatca 12300
gcccgtcata cttgaagcta gacaggctta tcttggaaca gaagaagatc gcttgccctc 12360
gcgcgcagat cagttggaag aatttgtcca ctacgtgaaa ggcgagatca ccaaggtagt 12420
cggcaataaa tgtctaaca ttcgttcaag ccgacgcgcg ttcggcgccg ggcttaactc 12480
aagcgttaga tgcactaagc acataattgc tcacagccaa actatcaggt caagtctgct 12540
tttattattt ttaagcgtgc ataataagcc ctacacaaat tgggagatat atcatgaaag 12600
gctggctttt tcttggtatc gcaatagttg gcgaagtaat cgcaacatcc gcattaaaat 12660
ctagcgaggg ctttactaag ctagcttgct tggctggtcc ggtaccgtga acgtcggtc 12720
gattgtacct gcgttcaaat actttcgat cgtgttcgcg gcttggccgg tgcgtcggt 12780
gatctcacgg atcgactgct tctctcgcaa cgccatccga cggatgatgt ttaaaagtcc 12840
catgtggatc actccgttgc cccgtcgctc accgtgttgg ggggaaggtg cacatggctc 12900
agttctcaat ggaaattatc tgcctaaccg gctcagttct cgttagaaac caacatgcaa 12960
gctccaccgg gtgcaaagcg gcagcgg 12987

```

<210> 25

<211> 13226

<212> DNA

<213> Artificial sequence

<220>

<223> T-DNA vector comprising a chimeric ParG expression reducing gene

<220>

```

<221> misc_feature
<222> (1)..(25)
<223> Left T-DNA border (C)

<220>
<221> misc_feature
<222> (58)..(318)
<223> 3' nos (C)

<220>
<221> misc_feature
<222> (337)..(888)
<223> bar coding region (C)

<220>
<221> misc_feature
<222> (889)..(1721)
<223> 35S3 promoter region (C)

<220>
<221> misc_feature
<222> (1778)..(3123)
<223> 35S promoter region

<220>
<221> misc_feature
<222> (3130)..(3431)
<223> part of ParG homologue of Zea mays

<220>
<221> misc_feature
<222> (3464)..(4205)
<223> Pdk-intron

<220>
<221> misc_feature
<222> (4238)..(4536)
<223> part of ParG homologue of Zea mays (C)

<220>
<221> misc_feature
<222> (4549)..(5259)
<223> 3' ocs

<220>
<221> misc_feature
<222> (5281)..(5305)
<223> Right T-dna border (C)

<400> 25
cggcaggata tattcaattg taaatggctc catggcgatc gctctagagg atcttcccga 60
tctagtaaca tagatgacac cgcgcgcgat aatttatcct agtttgcgcg ctatattttg 120
ttttctatcg cgtattaaat gtataattgc gggactctaa tcataaaaac ccatctcata 180
aataacgtca tgcattacat gttaattatt acatgcttaa cgtaattcaa cagaaattat 240
atgataatca tcgcaagacc ggcaacagga ttcaatctta agaaacttta ttgccaaatg 300
tttgaacgat ctgcttcgga tcctagacgc gtgagatcag atctcggtga cgggcaggac 360
cggacggggc ggtaccggca ggctgaagtc cagctgccag aaaccacgt catgccagtt 420
cccgtgcttg aagccggccg cccgcagcat gccgcggggg gcatatccga gcgcctcgtg 480
catgcgcacg ctcggttcgt tgggcagccc gatgacagcg accacgctct tgaagccctg 540
tgcctccagg gacttcagca ggtgggtgta gagcgtggag cccagtccec tccgctggtg 600
gcgggggggag acgtacacgg tcgactcggc cgtccagtcg taggcgttgc gtgccttcca 660

```

ggggcccgcg	taggcgatgc	cggcgacctc	gccgtccacc	tcggcgacga	gccagggata	720
gcgctccgcg	agacggacga	ggctgcctcg	ccactcctgc	ggttcctgcg	gctcggtagc	780
gaagtggacc	gtgcttgctc	cgatgtagtg	gttgacgatg	gtgcagaccg	ccggcatgtc	840
cgctccgggtg	gcacggcgga	tgctcgccgg	gcgtcgcttc	gggtccatgg	ttatagagag	900
agagatagat	ttatagagag	agactgggtga	tttcagcgtg	tcctctccaa	atgaaatgaa	960
cttccttata	tagaggaagg	gtcttgcgaa	ggatagtggg	attgtgcgtc	atcccttacg	1020
tcagtggaga	tgtcacatca	atccacttgc	tttgaagacg	tggttggaac	gtcttctttt	1080
tccacgatgc	tcctcgtggg	tgggggtcca	tctttgggac	cactgtcggc	agaggcatct	1140
tgaatgatag	cctttccttt	atcgcaatga	tggcatttgt	aggagccacc	ttccttttct	1200
actgtccttt	cgatgaagtg	acagatagct	gggcaatgga	atccgaggag	gtttcccgaa	1260
attatccttt	gttgaaaagt	ctcaatagcc	ctttgggtctt	ctgagactgt	atctttgaca	1320
tttttgaggt	agaccagagt	gtcgtgctcc	accatgttga	cgaagatttt	cttcttgta	1380
ttgagtcgta	aaagactctg	tatgaactgt	tcgccagtct	tcacggcgag	ttctgttaga	1440
tcctcgattt	gaatcttaga	ctccatgcat	ggccttagat	tcagttagaa	ctaccttttt	1500
agagactcca	atctctatta	cttgcccttg	tttatgaagc	aagccttgaa	tcgtccatac	1560
tggaatagta	cttctgatct	tgagaaatat	gtctttctct	gtgttcttga	tgcaattagt	1620
cctgaatctt	ttgactgcat	ctttaacctt	cttgggaagg	tatttgatct	cctggagatt	1680
gttactcggg	tagatcgtct	tgtagagacc	ttgtgcgtag	gaacgcggcc	gcgtatacgt	1740
atcgatatct	tcgaattcga	gtcgtcgag	cggcgctcg	acgaattaat	tccaatccca	1800
caaaaatctg	agcttaacag	cacagttgct	cctctcagag	cagaatcggg	tattcaacac	1860
cctcatatca	actactacgt	tgtgtataac	ggtcacatg	ccggtatata	cgatgactgg	1920
gggtgtacaa	aggcggcaac	aaacgcggtt	cccgaggtt	cacacaagaa	atttgccact	1980
attacagagg	caagagcagc	agctgacgcg	tacacaacaa	gtcagcaaac	agacaggttg	2040
aacttcatcc	ccaaaggaga	agctcaactc	aagcccaaga	gctttgctaa	ggccctaaca	2100
agcccaccaa	agcaaaaagc	ccactggctc	acgctaggaa	ccaaaaggcc	cagcagtgat	2160
ccagcccaaa	aagagatctc	ctttgccccg	gagattacaa	tggaagattt	cctctatctt	2220
tacgatctag	gaaggaagtt	cgaaggtgaa	ggtagcagaa	ctatgttcac	cactgataat	2280
gagaagggtta	gcctcttcaa	tttcagaaag	aatgctgacc	cacagatggt	tagagaggcc	2340
tacgcagcag	gtctcatcaa	gacgatctac	ccgagtaaca	atctccagga	gatcaaatac	2400
cttcccaaga	aggttaaaaga	tgcagtcaaa	agatttcagga	ctaattgcat	caagaacaca	2460
gagaaagaca	tatttctcaa	gatcagaagt	actattccag	tatggacgat	tcaaggcttg	2520
cttcataaac	caaggcaagt	aatagagatt	ggagtctcta	aaaaggtagt	tcctactgaa	2580
tctaaggcca	tgcatggagt	ctaagattca	aatcgaggat	ctaacagaa	tcgccgtgaa	2640
gactggcgaa	cagttcatac	agagtctttt	acgactcaat	gacaagaaga	aaatcttcgt	2700
caacatgggtg	gagcagcaga	ctctggtcta	ctccaaaaat	gtcaaagata	cagtctcaga	2760
agaccaaagg	gctatttgaga	cttttcaaca	aaggataatt	tcgggaaacc	tcctcggatt	2820
ccattgcccc	gctatctgtc	acttcatcga	aaggacagta	gaaaagggaag	gtggctccta	2880
caaagtccat	cattgcgata	aaggaaaggc	tatcattcaa	gatctctctg	ccgacagtgg	2940
tcccaaagat	ggacccccac	ccacgaggag	catcgtggaa	aaagaagacg	ttccaaccac	3000
gtcttcaaag	caagtggatt	gatgtgacat	ctccactgac	gtaagggatg	acgcacaatc	3060
ccactatcct	tcgcaagacc	cttctcttat	ataaggaagt	tcatttcatt	tggaaggagc	3120
acgctcgagg	aatctggctg	tctcctaagg	gaagtgaaca	aggcattttg	tggaattttc	3180
gatcaatcga	aacaccatct	ctatgcgaag	cttttccagg	atttgcacaa	caaggatgac	3240
ttttcaagca	tcaattccag	tgagtacgta	ggagtttcaa	caggaaactg	gggttggtgt	3300
gcttttggtg	gaaaccctga	aatcaagagc	atgattcagt	ggattgctgc	atcacaggct	3360
cttcgccctt	ttgttaatta	ctacactttt	gagaacgtgt	ctctgcaaag	attagaggag	3420
gtgatccagt	gggtacccca	gcttggtaag	gaaataatta	ttttcttttt	tccttttagt	3480
ataaaaatag	taagtgtatg	taattagtat	gattataata	atatagttgt	tataattgtg	3540
aaaaaataat	ttataaatat	attgtttaca	taaacaacat	agtaatgtaa	aaaaatatga	3600
caagtgtatg	gtaagacgaa	gaagataaaa	gttgagagta	agtataattat	ttttaatgaa	3660
tttgatcgaa	catgtaagat	gatatactag	cattaatatt	tgtttttaatc	ataatagttaa	3720
ttctagctgg	tttgatgaat	taaatatcaa	tgataaaaata	ctatagtaaa	aataagaata	3780
aataaattaa	ataaatattt	ttttatgatt	ataagtttat	tatataatta	aatatctata	3840
ccattactaa	atattttagt	ttaaaagtta	ataaatattt	tgttagaaat	tccaatctgc	3900
ttgtaattta	tcaataaaca	aaatattaaa	taacaagcta	aagtaacaaa	taatatcaaa	3960
ctaatagaaa	cagtaatcta	atgtaacaaa	acataatcta	atgctaata	aacaaagcgc	4020
aagatctatc	attttatata	gtattatttt	caatcaacat	tcttattaat	ttctaaataa	4080
tacttgtagt	tttattaact	tctaaatgga	ttgactatta	attaaatgaa	ttagtgcgaac	4140
atgaataaac	aaggtaacat	gatagatcat	gtcattgtgt	tatcattgat	cttacatttg	4200
gattgattac	agttgggaag	ctgggttcga	aatcgatcac	tggtatcacct	cctctaactct	4260
ttgcagagac	acgttctcaa	aagtgtagta	attaacaaaa	gggcgaagag	cctgtgatgc	4320

agcaatccac	tgaatcatgc	tcttgatttc	agggtttcca	ccaaaagcac	cacaacccca	4380
gtttcctggt	gaaactccta	cgtactcact	ggaattgatg	cttgaaaagt	catccttggt	4440
gtgcaaatcc	tggaaaagt	tcgcatagag	atgggtgttc	gattgatcga	aaaatccaca	4500
aaatgccttg	ttcacttccc	ttaggagaca	gccagattct	ctagagtcc	gctttaatga	4560
gatatgcgag	acgcctatga	tcgcatgata	tttgctttca	attctgttgt	gcacgttgta	4620
aaaaacctga	gcatgtgtag	ctcagatcct	taccgcccgt	ttcggttcat	tctaataaat	4680
atatcaccgc	ttactatcgt	atttttatga	ataatattct	ccgttcaatt	tactgattgt	4740
accctactac	ttatatgtac	aataattaaa	tgaatacaat	atattgtgct	gaatagggtt	4800
atagcgacat	ctatgataga	gcgccacaat	aacaaacaat	tgcgttttat	tattacaaat	4860
ccaattttta	aaaaagcggc	agaaccgggc	aaactaaaa	gactgattac	ataaatctta	4920
ttcaaatctt	aaaaggcccc	aggggctagt	atctacgaca	caccgagcgg	cgaactaata	4980
acgttcactg	aagggaactc	cgggtccccg	ccggcgcgca	tgggtgagat	tccttgaagt	5040
tgagtattgg	ccgtccgctc	taccgaaagt	tacgggcacc	attcaaccgc	gtccagcacg	5100
gcggccgggt	aaccgacttg	ctgccccgag	aattatgcag	catttttttg	gtgatgtgg	5160
gccccaaatg	aagtgcagggt	caaaccctga	cagtgcagac	aaatcggttg	gcgggtccag	5220
ggcgaatttt	gcgacaacat	gtcagggctc	agcaggacct	gcaggtcgac	ggccgagtac	5280
tggcaggata	tataccgttg	taatttgctg	cgtgtgaata	agtcgctgtg	tatgtttgtt	5340
tgattgtttc	tgttgagggtg	cagcccattt	caccggacaa	gtcggctaga	ttgatttagc	5400
cctgatgaac	tgcgaggggg	aagccatctt	gagcgcgaa	tgggaatgga	tttcgttgta	5460
caacgagacg	acagaacacc	cacgggaccg	agcttcgac	gagcatcaaa	tgaactgca	5520
atttattcat	atcaggatta	tcaataccat	atttttgaaa	aagccgtttc	tgtaataag	5580
gagaaaaactc	accgaggcag	ttccatagga	tggcaagatc	ctggtatcgg	tctgcgattc	5640
cgactcgtcc	aactcaata	caacctatta	atttccctc	gtcaaaaata	aggttatcaa	5700
gtgagaaatc	accatgagtg	acgactgaat	ccggtgagaa	tggcaaaagt	ttatgcattt	5760
ctttccagac	ttgttcaaca	ggccagccat	tacgctcgct	atcaaaatca	ctcgcatcaa	5820
ccaaaccgtt	attcattcgt	gattgcgcct	gagcgagacg	aaatacgccg	ctgttaaaag	5880
gacaattaca	aacaggaatc	gaatgcaacc	ggcgcaggaa	cactgccagc	gcatacaaa	5940
tattttcacc	tgaatcagga	tattcttcta	atacctggaa	tgctgttttt	ccggggatcg	6000
cagtggtag	taacctgca	tacggagata	taacggataa	atgcttgatg	gtcggaagag	6060
gcataaattc	cgtagccag	tttagtctga	ccatctcatc	tgtaacatca	ttggcaacgc	6120
tacctttgcc	atgtttcaga	aacaactctg	gcgcctcggg	cttcccatag	aatcgataga	6180
ttgtcgacc	tgattgccc	acattatccg	aatctggcaa	ttccggttcg	cttgctgtcc	6240
ataaaaccgc	ccagtctagc	tatcgccatg	taagcccact	gcaagctacc	tgctttctct	6300
ttgcgcttgc	gttttcggga	tcttcttgag	atcctttttt	tctgcgcgta	atctgctgct	6360
tgcaaacaaa	aaaaccaccg	ctaccagcgg	tggtttgttt	gccggatcaa	gagctaccaa	6420
ctctttttcc	gaaggtaact	ggcttcagca	gagcgcagat	accaaatact	gtccttctag	6480
tgtagccgta	gttaggccac	cacttcaaga	actctgtagc	accgcctaca	tacctcgctc	6540
tgctaatacct	gttaccagtg	gctgctgcca	gtggcgataa	gtcgtgtctt	accgggttgg	6600
actcaagacg	atagttaccg	gataaggcgc	agcggtcggg	ctgaacgggg	ggttcgtgca	6660
cacagcccag	cttgaggcga	acgacctaca	ccgaactgag	atacctacag	cgtgagctat	6720
gagaaaagcgc	cagccttccc	gaaggagaa	aggcggacag	gtatccggtg	agcggcaggg	6780
tcggaacagg	agagcgcacg	agggagcttc	caaggggaaa	cgcttggtat	ctttatagtc	6840
ctgtcgggtt	tcgccacctc	tgacttgagc	gtcgattttt	gtgatgtctg	tcaggggggc	6900
ggagcctatg	gaaaaacgcc	agcaacgcgg	ccttttttacg	gttcctggcc	ttttgctggc	6960
cttttgctca	catgttcttt	cctgcgttat	cccctgattc	tgtggataac	cgtattaccg	7020
cctttgagtg	agctgatacc	gctcgccgca	gccgaacgac	cgagcgcagc	gagtcagtga	7080
gcgaggaagc	ggaagagcgc	ctgatgcggg	attttctcct	tacgcatctg	tgcggtattt	7140
cacaccgcat	atggtgcact	ctcagtacaa	tctgctctga	tgcgcgatag	ttaagccagt	7200
atacactccg	ctatcgctac	gtgactgggt	catggctgcg	ccccgacacc	cgccaacacc	7260
cgctgacgcg	ccctgacggg	cttgtctgct	cccggcatcc	gcttacagac	aagctgtgac	7320
cgtctccggg	agctgcatgt	gtcagagggt	ttcaccgtca	tcaccgaac	gcgcgaggca	7380
gggtgccttg	atgtggggcg	cggcggtcga	gtggcgacgg	cgcggcttgt	cgcgcctctg	7440
gtagattgcc	tggccttagg	ccagccattt	ttgagcggcc	agcggccgcg	ataggccgac	7500
gcgaagcggc	ggggcgtagg	gagcgcagcg	accgaagggt	aggcgttttt	tgcagctctt	7560
cggctgtgcg	ctggccagac	agttatgcac	aggccaggcg	ggttttaaga	gttttaataa	7620
gttttaaaga	gttttaggcg	gaaaaatcgc	cttttttctc	ttttatatca	gtcacttaca	7680
tgtgtgaccg	gttcccattg	tacggctttg	ggttcccatt	gtacgggttc	cggttcccaa	7740
tgtaaggctt	tgggttccca	atgtacgtgc	tatccacagg	aaagagacct	tttcgacctt	7800
tttccctcgc	tagggcaatt	tgccctagca	tctgctccgt	acattaggaa	ccggcggtatg	7860
cttcgcccctc	gatacaggtg	cggtagcgca	tgactaggat	cgggcccagcc	tgccccgcct	7920
cctccttcaa	atcgtactcc	ggcaggtcat	ttgacccgat	cagcttgccg	acggtgaaac	7980

agaactttctt	gaactctccg	gcgctgccac	tgcgttcgta	gatcgtcttg	aacaaccatc	8040
tggcttctgc	cttgccctgc	gcgcggcggtg	ccaggcggtg	gagaaaacgg	ccgatgccgg	8100
gatcgatcaa	aaagtaactc	gggtgaaccg	tcagcacgtc	cgggttcttg	ccttctgtga	8160
tctcgcggta	catccaatca	gctagctcga	tctcgatgta	ctccggccgc	ccggtttcgc	8220
tctttacgat	cttgtagcgg	ctaatacagg	cttcaccctc	ggataccgtc	accaggcggc	8280
cgttcttgge	cttcttcgta	cgcgtcatgg	caacgtgcgt	ggtgtttaac	cgaatgcagg	8340
tttctaccag	gtcgtctttc	tgttttccgc	catcggctcg	ccggcagaac	ttgagtacgt	8400
ccgcaacgtg	tggacggaac	acgcggccgg	gcttgtctcc	cttcccttcc	cggtatcggt	8460
tcattgattc	ggttagatgg	gaaaccgcca	tcagtaccag	gtcgtaatcc	cacacactgg	8520
ccatgccggc	cggccctgcg	gaaacctcta	cgtgcccgct	tggagctcgc	tagcggatca	8580
cctcggcagc	tcgtcgggtc	cgtctcgaca	gacggaaaac	ggccacgtcc	atgatgctgc	8640
gagtatcgcg	ggtgccccag	tcatagagca	gtgcctgggt	aaaatctggg	tgctcgtcgc	8700
ccttggggcg	cttcctaata	gacggcgcac	cggctgccgg	cggttgccgg	gattctttgc	8760
ggattcgatc	agcggccgct	tgccacgatt	caccggggcg	tgcttctgcc	tcgatgcgtt	8820
gccgctgggc	ggcctgcgcg	gccttcaact	tctccaccag	gtcatcaccc	agcgccgcgc	8880
cgatttgatc	cgggcccggg	ggtttgcgac	cgtcacgcgc	attcctcggg	cttggggggt	8940
ccagtgccat	tgaggggccg	gcagacaacc	cagccgctta	cgcctggcca	accgcccgtt	9000
cctccacaca	tggggcacgc	cacggcgctc	gtgcctgggt	gttcttgatt	ttccatgccg	9060
cctcctttag	ccgctaaaat	tcattctactc	atttattcat	ttgctcattt	actctggtag	9120
ctgcgcgatg	tattcagata	gcagctcggg	aatggtcttg	ccttggcgta	ccgcgtacat	9180
cttcagcttg	gtgtgatcct	ccgccggcaa	ctgaaagtgt	accgccttca	tggttgccgt	9240
gtctgccagg	ctggccaacg	ttgcagcctt	gctgtgcgt	gcgctcggac	ggccggcact	9300
tagcgtgttt	gtgcttttgc	tcattttctc	tttacctcat	taactcaaat	gagttttgat	9360
ttaatttcag	cggccagcgc	ctggacctcg	cggccagcgt	cgcctcggg	ttctgattca	9420
agaacgggtg	tgccggcggc	ggcagtgccct	gggtagctca	cgcgctgcgt	gatacgggac	9480
tcaagaatgg	gcagctcgta	cccgccagc	gcctcggcaa	cctcaccgcc	gatgcgcgtg	9540
cctttgatcg	cccgcgacac	gacaaaggcc	gcttgtagcc	ttccatccgt	gacctcaatg	9600
cgtgtcttaa	ccagctccac	caggtcggcg	gtggcccata	tgtcgtaagg	gcttggctgc	9660
accggaatca	gcacgaagtc	ggctgccttg	atcgcggaca	cagccaagtc	cgcgcgctgg	9720
ggcgctccgt	cgatcactac	gaagtcgcgc	cggccgatgg	ccttcacgtc	gcggtcaatc	9780
gtcggggcgt	cgatgccgac	aacggttagc	gggtgatctt	cccgcacggc	cgcccaatcg	9840
cgggcactgc	cctggggatc	ggaatcgact	aacagaacat	cggccccggc	gagttgcagg	9900
gcgcgggcta	gatgggttgc	gatggctcgt	ttgcctgacc	cgcctttctg	gttaagtaca	9960
gcgataacct	tcattgcgttc	cccttgcgta	tttgtttatt	tactcatcgc	atcatatacg	10020
cagcgaccgc	atgacgcaag	ctgttttact	caaatacaca	tcaccttttt	agacggcggc	10080
gctcgggttc	ttcagcggcg	aagctggccg	cgaggccgc	cagcttgcca	tcagacaaac	10140
cggccaggat	ttcatgcagc	cgcacgggtg	agacgtgcgc	gggcggctcg	aacacgtacc	10200
cggccgcgat	catctccgcc	tcgatctctt	cggtaatgaa	aaacggttcg	tcctggccgt	10260
cctggtgcgg	tttcatgett	gttcctcttg	gcgttcattc	tcggcgccgc	ccaggggcgtc	10320
ggcctcggtc	aatgcgtcct	cacggaaggc	accgcgcgc	ctggcctcgg	tgggcgctcac	10380
ttcctcgtcg	cgtcaagtg	cgcgggtacg	ggtcgagcga	tgacgcgcaa	gcagtgcagc	10440
cgcctctttc	acggtgcggc	cttcctggtc	gatcagctcg	cgggcgtgcg	cgatctgtgc	10500
cggggtgagg	gtaggggcgg	ggccaaactt	cacgcctcgg	gccttggcgg	cctcgcgccc	10560
gctccgggtg	cggtcgatga	ttagggaacg	ctcgaaactc	gcaatgccgg	cgaacacggt	10620
caacaccatg	cggccggccg	gcgtgggtgt	gtcggcccac	ggctctgcc	ggctacgcag	10680
gcccgcgcgc	gcctcctgga	tgcgctcggc	aatgtccagt	aggtcgcggg	tgctgcgggc	10740
caggcggtct	agcctgggtc	gtgcacaaac	gtgcgaggg	cgtaggtggg	caagcatcct	10800
ggccagctcc	ggcggtcgc	ccctgggtgc	gggtatcttc	tcggaaaaca	gcttgggtgca	10860
gccggccgcg	tgagttcgg	cccgttggtt	ggtaagtc	tggtcgtcgg	tgctgacgcg	10920
ggcatagccc	agcaggccag	cggcggcgct	cttggtcatg	gcgtaatgtc	tcgggttcta	10980
gtcgcaagta	ttctacttta	tgcgactaaa	acacgcgaca	agaaaacgcc	aggaaaagg	11040
caggggcgga	gcctgtcgcg	taacttagga	cttgtgcgac	atgtcgtttt	cagaagacgg	11100
ctgcactgaa	cgtcagaagc	cgactgcact	atagcagcgg	aggggttggg	tcgatccctg	11160
ctcgcgcagg	ctgggtgcc	agctctcggg	taacatcaag	gcccgatcct	tggaagccctt	11220
gccctccgcg	acgatgatcg	tgcggtgatc	gaaatccaga	tccttgaccc	gcagttgcaa	11280
accctcactg	atccgcgatg	ccgttccata	cagaagctgg	gcgaacaaac	gatgctcgcc	11340
ttccagaaaa	ccgaggatgc	gaaccacttc	atccgggggtc	agcaccaccg	gcaagcgcgc	11400
ggacggccga	ggtcttccga	tctcctgaag	ccagggcaga	tcggtgcaca	gcacttgccg	11460
tagaagaaca	gcaaggccgc	caatgcctga	cgatgcgtgg	agaccgaaac	cttgcgctcg	11520
ttcgccagcc	aggacagaaa	tgctcgcact	tcgctgctgc	ccaaggttgc	cgggtgacgc	11580
acaccgtgga	aacgggatgaa	ggcacgaacc	cagtgagcat	aagcctgttc	ggttcgtaag	11640

ctgtaatgca	agtagcgat	gcgctcacgc	aactggtcca	gaaccttgac	cgaacgcagc	11700
ggtggtaacg	gcgcagtgge	ggttttcatg	gcttggtatg	actgtttttt	tggggtagacg	11760
tctatgcctc	gggcatcaca	gcagcaagcg	cgttacgcgc	tgggtcgatg	tttcatgtta	11820
tggagcagca	acgatgttac	gcagcagggc	agtcgcccta	aaacaaagtt	aaacatcatg	11880
agggaaagcgg	tgatcgccga	agtatcgact	caactatcag	aggtagttgg	cgatcatcgag	11940
cgccatctcg	aaccgcagtt	gctggccgta	catttgtagc	gctccgcagt	ggatggcggc	12000
ctgaagccac	acagtgatat	tgatttgctg	gttacgggtg	ccgtaaggct	tgatgaaaca	12060
acgcggcgag	ctttgatcaa	cgaccttttg	gaaacttcgg	cttcccctgg	agagagcgag	12120
attctccgcg	ctgtagaagt	caccattgtt	gtgcacgacg	acatcattcc	gtggcggtat	12180
ccagctaagc	gcgaactgca	atltggagaa	tggcagcgca	atgacattct	tgcaggatc	12240
ttcgagccag	ccacgatcga	cattgatctg	gctatcttgc	tgacaaaagc	aagagaacat	12300
agcggtgcct	tggtaggtcc	agcggcggag	gaactctttg	atccggttcc	tgaacaggat	12360
ctatttgagg	cgctaaatga	aaccttaacg	ctatggaact	cgccgcccca	ctgggctggc	12420
gatgagcgaa	atgtagtgtc	tacgttggtc	cgcatttggt	acagcgagc	aaccggcaaa	12480
atcgcgccga	aggatgtcgc	tgccgactgg	gcaatggagc	gcctgccggc	ccagtatcag	12540
cccgatcata	ttgaagctag	acaggcttat	cttggaacag	aagaagatcg	cttggcctcg	12600
cgcgagatc	agttggaaga	atltgtccac	tacgtgaaag	gcgagatcac	caaggtagtc	12660
ggcaaataat	gtctaacaat	tcgttcaagc	cgacgccgct	tcgcggcgcg	gcttaactca	12720
agcgtagat	gcactaagca	cataattgct	cacagccaaa	ctatcaggtc	aagtctgctt	12780
ttattatatt	taagcgtgca	taataagccc	tacacaaatt	gggagatata	tcatgaaagg	12840
ctggcttttt	cttggtatcg	caatagttgg	cgaagtaatc	gcaacatccg	cattaaaatc	12900
tagcgagggc	ttactaagc	tagcttgctt	ggtcggtccg	gtaccgtgaa	cgtcggctcg	12960
attgtacctg	cgttcaaata	ctttgcgac	gtgttgccgc	cctgcccggt	gcgtcggtcg	13020
atctcacgga	tcgactgctt	ctctcgcaac	gccatccgac	ggatgatgtt	taaaagtccc	13080
atgtggatca	ctccgttgcc	ccgtcgctca	ccgtgttggg	gggaagggtc	acatggctca	13140
gttctcaatg	gaaattatct	gcctaaccgg	ctcagttctg	cgtagaaacc	aacatgcaag	13200
ctccaccggg	tgcaagcg	cagcgg				13226